

Cytotoxicity of airborne particulate matter (PM₁₀) from dust storm and inversion conditions assessed by MTT assay

Maryam Faraji¹, Ramin Nabizadeh Nodehi^{1,2}, Kazem Naddafi^{1,2}, Zahra Pourpak³, Zahra Alizadeh³, Soheila Rezaei⁴, Alireza Mesdaghinia^{1,2,*}

¹ Department of Environmental Health Engineering, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

² Center for Air Pollution Research (CAPR), Institute for Environmental Research (IER), Tehran University of Medical Sciences, Tehran, Iran

³ Immunology, Asthma and Allergy Research Institute, Tehran University of Medical Sciences, Tehran, Iran

⁴ Social Determinants of Health Research Center, Yasuj University of Medical Sciences, Yasuj, Iran

ARTICLE INFORMATION

Article Chronology:

Received 27 July 2018

Revised 26 August 2018

Accepted 9 September 2018

Published 29 September 2018

Keywords:

Toxicological studies; Dust storm; Air pollution; MTT test; Cytotoxicity; PBMCs

CORRESPONDING AUTHOR:

mesdaghinia@sina.tums.ac.ir

Tel: (+98 21 88954914

Fax: (+98 21) 88950188

ABSTRACT:

Introduction: Airborne particles generate acute and chronic toxic effects on the human health. Cytotoxicity of air pollutants can be investigated through cytotoxicity assays. In this study, cytotoxicity of PM₁₀ (particles ≤ 10 μm in diameter) from dust storm and inversion condition was compared through MTT assay on the human peripheral blood mononuclear cells (PBMCs) in vitro.

Materials and methods: PM₁₀ was sampled in Tehran, Iran, 2016, in dust storm and inversion. PBMCs were isolated from the whole blood sample through Ficoll - Hypaque gradient method. Cells were treated with two suspensions of the PM₁₀ from dust storm and inversion at different concentrations (50, 100, 150, 200, 250, 300, 350 and 400 μg / mL) for 24 h. Cell viability was assessed by MTT test and reported in respect to the viability in untreated cells as negative control.

Results: During the sampling period, June 6 and 12 - 15 November, 2016, were selected as the dusty and inversion days, respectively. Daily average PM₁₀ in dust storm and inversion conditions were found of 220 and 345 μg / m³, respectively. Mean of viability in the PBMCs treated by the samples from dust storm and inversion was found 85.79 ± 9.97 % and 81.58 ± 11.72%, respectively. The cell viability values were obtained between 78 - 96 % for PM₁₀ related dust storm condition and 70 - 92 % for PM₁₀ sampled in inversion days.

Conclusion: The results showed that the PM₁₀ from dust storm as well as from inversion had the cytotoxicity effects on PBMCs. The particles related to the inversion caused toxic effects more than those from dust storm at all concentrations.

Introduction

Air pollution is a mixture of different pollutants included particulate matter (PM) and gases. PM can be originated from the anthropogenic and natural

sources. Automobiles, power generators, industries, waste incinerators, biomass burning and domestic heating and cooking are the chief anthropogenic sources [1, 2]. Dust storm is one of the

natural sources, which it has mainly participated in the air pollution of several countries in the Middle East [3, 4]. Air particles can cause toxic health effects on the human such as lung cancer and leukemia [5], allergic reactions, asthma [6], eye infections, meningitis [7, 8], mortality and hospitalization due to the cardiovascular and respiratory diseases, chronic obstructive pulmonary disease, acute myocardial infarction [9] and infertility [10]. The International Agency for Research on Cancer (IARC) has considered the outdoor air pollution and PM as the definite human carcinogen, which mainly cause the lung cancer [11].

Cytotoxicity of chemicals can be determined through *in vitro* cytotoxicity assays. Their performance is based on the several cell functions such as enzyme activity, membrane permeability and adherence of cell, production of adenosine triphosphate (ATP) and co-enzymes and the activity of nucleotide uptake. According to these functions, various methods are used for measuring the cytotoxicity such as crystal violet, colony formation, tritium-labeled thymidine uptake and colorimetric method by using tetrazolium dye, briefly MTT. Among mentioned methods, MTT is safe, easy-to-use, reliable with a high reproducibility. In MTT assay, NAD(P)H-dependent cellular oxidoreductase enzymes in the active cells reduce the tetrazolium dye MTT (3-(4, 5-dimethylthiazolyl-2)-2,5-diphenyltetrazolium bromide) to the intracellular purple formazan [12].

MTT method was studied to assay cytotoxicity of $PM_{2.5}$ (particles $\leq 2.5 \mu\text{m}$ in diameter) and $PM_{2.5-10}$ (particles between 2.5-10 μm in diameter) ambient air pollutants [13], dust storm $PM_{2.5}$ [14], urban PM_{10} and $PM_{2.5}$ [15], Fine particulate matter (PM_1 and $PM_{2.5}$) in the Milan urban area [16], ambient $PM_{2.5}$ [17].

To the best of our knowledge, toxicity of PM_{10} (particles $\leq 10 \mu\text{m}$ in diameter) from dust storm and inversion sources have not yet been com-

pared by MTT method. So, this study aimed to compare cytotoxicity of PM_{10} collected in dust storm and inversion conditions through MTT assay on the human peripheral blood mononuclear cells (PBMCs).

Materials and methods

The study stages, including sampling and preparation of PM_{10} , PBMCs isolation, cell treatment with the particle suspension, MTT assay and statistical analysis, are visualized in Fig. 1. Chemical analysis on the samples has been reported in other study [3].

Air sampling

Particles were collected in Tehran, Iran ($35^{\circ}70'66.00''\text{N}$, $51^{\circ}39'38.55''\text{E}$) (Fig. 2). Since, dust storm and inversion occur in Tehran mainly in spring and autumn, respectively, 24 h sampling was done over the two periods, from April 26 to June 7 and September 24 to November 15, 2016.




Fig. 1. Study stages

Particles were sampled using a high-volume sampler (1.3 – 1.7 m³ / min) (Grasebey, USA) equipped with fiberglass filter (8×10 inch, grade G 653 Whatman, USA). Filters were weighed with an analytical balance (± 10 mg) before and after sampling to calculate the mass of sampled PM. The sampler was installed based on the United States Environmental Protection Agency (USEPA) instruction [18] at the height of 10 m above the ground level far away from any obstruction to prevent the potential effects of natural and anthropogenic obstacles on the air flow and PM concentration.

Preparation of PM₁₀ samples

In the first stage of PM₁₀ preparation, two condi-

tions of dust storm and inversion must be differentiated. Dust storm and inversion conditions were distinguished according to the Hoffmann's criteria [19] (Table 1) and the report of the Tehran Air Quality Control Company (TAQCC), respectively. The particles were extracted from the filters through the dry ultrasonic (Elma-ultrasonic, Germany) followed by sweeping with a smooth brush [3]. The extracted PM₁₀ was weighed and stored into the endotoxin - free vials at -18°C until their use in biological test. Since, the fiberglass fibers have toxicity effects on cells [15], their presence was inspected in the extracted samples by using a scanning electron microscope (SEM) (HITACHI, SU3500, Japan)




Fig. 2. Map of study area and PM₁₀ sampling station

Table 1. Dust storm classification

Category	Visibility (m)	Wind speed (m / s)	PM ₁₀ (µg / m ³ .h)
Dusty Air (DA)	Haze	-	50-200
Light Dust Storm (DS1)	< 2000	-	200 - 500
Dust Storm (DS2)	< 1000	> 17	500 - 2000
Strong Dust Storm (DS3)	< 200	> 20	2000 - 5000
Serious Strong DS (DS4)	< 50	> 25	> 5000

PBMCs isolation and in vitro treatment

The volume of 20 mL whole blood sample was collected from the healthy volunteer and put into the heparinized tube as the anticoagulant and processed within 2 h. The method of Ficoll-Hypaque gradient was used to isolate the PBMCs. Briefly, 40 mL Ca²⁺/Mg²⁺ -free PBS (Biosera, France) was added to whole blood sample in the laminar flux hood to dilute it. Cells were isolated from the diluted blood with addition of 30 mL Ficoll-Hypaque solution (Biosera, France) by density centrifugation (22 min, 2000 rpm, no acceleration, no brake). Then, layer of PBMCs was collected and washed by the lysis buffer and isolation buffer and isolated through centrifugation (400 g, 14 min, acceleration 6, brake 4). The number of 200,000 cells were seeded in each well of a 96-well plate in 100 mL complete RPMI-1640 culture medium (Gibco BRL, San Diego, CA) and cultured in a humidified incubator at 37 °C with 5% (v/v) CO₂.

Two suspensions from dust storm and inversion were separately prepared in the culture medium. The cultured cells were treated with mentioned suspensions at different concentrations (50, 100, 150, 200, 250, 300, 350 and 400 µg / mL) and incubated at 37 °C for 24 h. Experiments at six concentrations were done in triplicate.

MTT assay

MTT solution in the final concentration of 0.5 mg / mL (Sigma Chemical Company, St. Louis, MO, USA) was added to the wells and incubated for

4 h. Insoluble formazan crystals generated as the byproduct of MTT assay were dissolved in 150 µL dimethylsulfoxide (DMSO) (Sigma Chemical Company, St. Louis, MO, USA). Then, the absorbance of samples was measured using a microplate reader (Bio-Tek Instruments, Winooski, VT, USA) at 570 nm. Finally, cell viability percent was calculated using Eq. (1):

(1)

$$\text{Cell viability}(\%) = (A_S - A_b) / (A_{NC} - A_b) \times 100$$

where A is the absorbance, b is the blank, and S and NC are the sample and negative control (untreated cells), respectively [20].

Statistical analysis

The data were analyzed with Excel 2016 software and reported as means ± standard deviation of three independent experiments.

Results and discussion

PM₁₀ concentration

According to the considered criteria, June 6 and November 12-15, 2016, had respectively the condition of dust storm and inversion over the particle sampling period. Particle sampling information at the two conditions is described in Table 2.

Particle extraction

Morphological structure of blank filter and extracted sample were shown in Fig. 3a and 3b, respectively.

Table 2. Information on the PM₁₀ sampling during dust storm and inversion conditions

Condition	Date	PM ₁₀ mass (g)	Air volume (m ³)	Daily average PM ₁₀ (µg / m ³)
Dust storm	June 6	0.57	1634.32	348.40
Inversion	November 12	0.48	2028.32	236.60
	November 13	0.41	2051.22	201.20
	November 14	0.43	2031.98	211.32
	November 15	0.43	1840.35	233.05




Fig. 3. Morphological structure of blank filter (a) and extracted sample (b)

Table 3. Viability percent of peripheral blood mononuclear cells (PBMCs) in MTT assay treated by PM₁₀ from dust storm and inversion at concentrations of 50 - 400 µg / mL.

	PM ₁₀ concentration (µg / ml)								Total
	50	100	150	200	250	300	350	400	
%Cell viability related to dust storm PM ₁₀									
Mean	96.00	93.33	92.33	90.33	87.33	83.00	78.33	65.67	85.79
SD*	4.36	5.69	5.51	10.02	8.62	5.29	3.21	5.03	9.97
% Cell viability related to inversion PM ₁₀									
Mean	92.00	90.00	91.00	87.00	84.33	79.33	70.33	58.67	81.58
SD*	4.36	4.00	6.56	3.00	5.03	4.04	5.51	3.21	11.72

*: Standard deviation

Cytotoxicity analysis

Mean of viability for PBMCs treated by the samples from dust storm and inversion was found $85.79 \pm 9.97\%$ and $81.58 \pm 11.72\%$, respectively. Viability percent of PBMCs at concentrations of 50 - 400 µg / mL represents in Table 3.

The dusty day with daily average PM₁₀ concentration of $348.40 \mu\text{g} / \text{m}^3$ could be categorized in the light dust storm class based on the Hoffmann classification in Table 1. Also, the world meteorological organization (WMO) classified dust events according to visibility into the four class-

es: (1) dust – in - suspension: visibility usually less than 10 km; (2) blowing dust: visibility 1-10 km; (3) dust storm: visibility 200 - 1000 m; and (4) severe dust storm: visibility less than 200 m [21]. In the current study, the dust storm day with the visibility of 5.95 km (Table 2) could be classified in the dust-in-suspension or blowing dust class. At both the conditions, daily average PM_{10} concentration was up to 6.96 times higher for the dust storm condition and 4.4 times higher for the inversion, as compared to that with the national guideline ($50 \mu\text{g} / \text{m}^3$).

Based on the comparison between the morphological structure of blank filter in Fig. 3a and the extracted particles in Fig. 3b, it can be concluded that fibers were not observed in the extracted particles. As a result, the observed toxicity effects in MTT assay could be certainly assigned to the particles. According to Table 3, it can be concluded that both types of particles had cytotoxicity effects on the PBMCs and they could suppress cell activity. The particles related to the inversion caused toxic effects more than those from dust storm at all concentrations (mean of viability of $85.79 \pm 9.97 \%$ in dust storm vs. $81.58 \pm 11.72 \%$ for inversion). The cell viability values were obtained between 78 – 96 % for PM_{10} related dust storm condition and 70 – 92 % for PM_{10} sampled from inversion. Cell viability was decreased less than 80 % at concentration of $350 \mu\text{g} / \text{mL}$ for the both types of particles (78 % for the PM from dust storm and 70 % for it from inversion). This reduced cell viability could be associated to the different physiochemical characteristics of particles in dust storm condition and inversion [18, 22]. Particles from dust storm and inversion had dissimilar nature because they release from different sources. Particles related to dust storm condition mainly disseminate from natural sources. Against, particles allocated to inversion in Tehran release from anthropogenic

sources mainly fossil fuels and vehicles [18]. In comparison between the chemical compositions, frequent ionic components were found NO_3^- , Cl^- , SO_4^{2-} and Ca^{+2} during dust storm (72.18 % of PM mass) and SO_4^{2-} , NO_3^- , NH_4^+ and Cl^- during inversion (87.78 % of PM mass). For both conditions, Si, Fe and Al were identified as the dominant elements respectively in dust storm condition 96.62 % of PM mass and in inversion 59.16 % of PM mass [3]. Usually, the particulates resulted from the anthropogenic sources have the higher concentrations of pollutants such as heavy metals and polyaromatic hydrocarbons [11].

Conclusion

In the present study, toxicology effect of PM_{10} was investigated on the PBMCs by MTT method. This effect was compared between particles from natural and anthropogenic sources, respectively dust storm and inversion. Results were verified that both category of the particles could decrease cell viability. But, particles from inversion condition generate the toxic effects more than those from dust storm. Different percentages of viability in the treated cells can be due to diverse chemical composition and physicochemical properties of the particles in studied conditions.

Financial supports

This study was financially supported by the Institute for Environmental Research (IER) of Tehran University of Medical Sciences (TUMS) (grant number 95-03-46-32844) and Iran National Science Foundation (INSF) (grant number 95831261).

Competing interests

All of the authors declare that they have no actual or potential personal or financial competing interests.

Acknowledgements

The authors would like to thank the IER and INSF for financially supports. Also, they are grateful to the Immunology, Asthma and Allergy Institute (IAARI) for technically support and staff collaboration.

Ethical considerations

This study was permitted by the ethics committee of the Tehran University of Medical Sciences. Written informed consent was obtained from volunteer before starting the study. Also, the results of the research will be published with the permission of the funders. Also, all authors agree to submit their manuscript to JAPH. The authors confirm that the manuscript have not been submitted or published elsewhere in any language.

References

- Hajizadeh Y, Mokhtari M, Faraji M, Mohammadi A, Nemati S, Ghanbari R, et al. Trends of BTEX in the central urban area of Iran: A preliminary study of photochemical ozone pollution and health risk assessment. *Atmospheric Pollution Research*. 2018;9:220-9.
- Nabizadeh R, Atafar Z, Faraji M. Spreadsheet model to design of hazardous waste incinerators. *Journal of Air Pollution and Health*. 2017;1(4):269-80.
- Faraji M, Pourpak Z, Naddafi K, Nodehi RN, Nicknam MH, Shamsipour M, et al. Effects of airborne particulate matter (PM10) from dust storm and thermal inversion on global DNA methylation in human peripheral blood mononuclear cells (PBMCs) in vitro. *Atmospheric Environment*. 2018;195: 170-8.
- Jaafari J, Naddafi K, Yunesian M, Nabizadeh R, Hassanvand MS, Ghoskhal MG, et al. Study of PM10, PM2.5, and PM1 levels in during dust storms and local air pollution events in urban and rural sites in Tehran. *Human and Ecological Risk Assessment: An International Journal*. 2018;24(2):482-93.
- Dehghani M, Keshtgar L, Javaheri MR, Derakhshan Z, Oliveri Conti G, Zuccarello P, et al. The effects of air pollutants on the mortality rate of lung cancer and leukemia. *Mol Med Rep*. 2017;15(5):3390-7.
- Ghoskhal MG, Ansarin K, Naddafi K, Nodehi RN, Yaghmaeian K, Hassanvand MS, et al. Short-term effects of particle size fractions on lung function of late adolescents. *Environmental Science and Pollution Research*. 2018:1-11.
- Ghoskhal MG, Ansarin K, Naddafi K, Nodehi RN, Yaghmaeian K, Hassanvand MS, et al. Prevalence of asthma and associated factors among male late adolescents in Tabriz, Iran. *Environmental Science and Pollution Research*. 2018;25(3):2184-93.
- Naddafi K, Atafar Z, Faraji M, Ghanbarian M, Rezaei S, Ghoskhal MG, et al. Health Effects of Airborne Particulate Matters (PM₁₀) during Dust Storm and Non-Dust Storm Conditions in Tehran. *Journal of Air Pollution and Health*. 2017;1(4):259-68.
- Miri M, Derakhshan Z, Allahabadi A, Ahmadi E, Oliveri Conti G, Ferrante M, et al. Mortality and morbidity due to exposure to outdoor air pollution in Mashhad metropolis, Iran. The AirQ model approach. *Environmental research*. 2016;151:451-7.
- Conti GO, Calogero AE, Giaccone F, Fiore M, Barchitta M, Agodi A, et al. B(a)P adduct levels and fertility: A cross-sectional study in a Sicilian population. *Molecular Medicine Reports*. 2017;15(5):3398-404.
- Ghanbarian M, Nicknam MH, Mesdaghinia A, Yunesian M, Hassanvand MS, Soleimanifar N, et al. Investigation and Comparison of In Vitro Genotoxic Potency of PM10 Collected in Rural and Urban Sites at Tehran in Different Metrological Conditions and Different Seasons. *Biological trace element research*. 2018:1-10.
- de Oliveira Galvão MF, de Oliveira Alves N, Ferreira PA, Caumo S, de Castro Vasconcellos P, Artaxo P, et al. Biomass burning particles in the Brazilian Amazon region: Mutagenic effects of nitro and oxy-PAHs and assessment of health risks. *Environmental Pollution*. 2018;233:960-70.
- Hsiao WW, Mo Z-Y, Fang M, Shi X-m, Wang F. Cytotoxicity of PM2.5 and PM2.5-10 ambient air pollutants assessed by the MTT and the Comet assays. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*. 2000;471(1):45-55.
- Meng Z, Zhang Q. Damage effects of dust storm PM2.5 on DNA in alveolar macrophages and lung cells of rats. *Food and chemical toxicology*. 2007;45(8):1368-74.
- Alfaro-Moreno E, Torres V, Miranda J, Martínez L, García-Cuellar C, Nawrot TS, et al. Induction of IL-6 and inhibition of IL-8 secretion in the human airway cell line Calu-3 by urban particulate matter collected with a modified method of PM sampling. *Environmental research*. 2009;109(5):528-35.
- Perrone MG, Gualtieri M, Ferrero L, Porto CL, Udisti R, Bolzacchini E, et al. Seasonal variations in chemical composition and in vitro biological effects of fine PM from Milan. *Chemosphere*. 2010;78(11):1368-77.
- MohseniBandpi A, Eslami A, Shahsavani A, Khodagholi F, Alinejad A. Physicochemical characterization of ambient PM 2.5 in Tehran air and its potential cytotoxicity in human lung epithelial cells (A549). *Science of The Total Environment*. 2017;593:182-90.
- Rezaei S, Naddafi K, Hassanvand MS, Nabizadeh R, Yunesian M, Ghanbarian M, et al. Physicochemical characteristics and oxidative potential of ambient air particulate matter (PM₁₀) during dust and non-dust storm

- events: a case study in Tehran, Iran. *Journal of Environmental Health Science and Engineering*. 2018;Online publish.
19. Hoffmann C, Funk R, Sommer M, Li Y. Temporal variations in PM₁₀ and particle size distribution during Asian dust storms in Inner Mongolia. *Atmospheric Environment*. 2008;42(36):8422-31.
 20. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of immunological methods*. 1983;65(1-2):55-63.
 21. Shao Y, Dong C. A review on East Asian dust storm climate, modelling and monitoring. *Global and Planetary Change*. 2006;52(1):1-22.
 22. Naimabadi A, Ghadiri A, Idani E, Babaei AA, Alavi N, Shirmardi M, et al. Chemical composition of PM₁₀ and its in vitro toxicological impacts on lung cells during the Middle Eastern Dust (MED) storms in Ahvaz, Iran. *Environmental Pollution*. 2016;211:316-24.