

## Health effects of airborne particulate matter related to traffic in Urmia, north-west Iran

Amir Mohammadi<sup>1,\*</sup>, Maryam Faraji<sup>2,3</sup>, Saeed Mousavi<sup>4</sup>, Sepideh Nemati<sup>5</sup>, Mojtaba Momtaz<sup>6</sup>, Ali Abdolahnejad<sup>6</sup>, Mohammad Miri<sup>7</sup>

<sup>1</sup> Department of Public Health, Maragheh University of Medical Sciences, Maragheh, Iran

<sup>2</sup> Environmental Health Engineering Research Center, Kerman University of Medical Sciences, Kerman, Iran

<sup>3</sup> Department of Environmental Health, School of Public Health, Kerman University of Medical Sciences, Kerman, Iran

<sup>4</sup> Environmental Research Center of West Azerbaijan, Urmia, Iran

<sup>5</sup> Health and Environment Research Center, Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

<sup>6</sup> Student Research Committee, Department of Environmental Health Engineering, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

<sup>7</sup> Department of Environmental Health Engineering, School of Health, Sabzevar University of Medical Sciences, Sabzevar, Iran

### ARTICLE INFORMATION

*Article Chronology:*

Received 29 April 2019

Revised 18 May 2019

Accepted 15 June 2019

Published 29 June 2019

*Keywords:*

Urmia city; Air pollution; Urmia lake; Health effects

### CORRESPONDING AUTHOR:

mohammadiurm@gmail.com

Tel: (+98 41) 37276363

Fax: (+98 41) 37276363

### ABSTRACT:

**Introduction:** This study aimed to evaluate exposure to PM<sub>10</sub> related to traffic and drying Lake Urmia and its effects on human health in north-west of Iran.

**Materials and methods:** Sampling and analysis was done during 2015 to 2016 using AirQ 2.2.3.

**Results:** The annual mean of PM<sub>10</sub> concentrations were 106 µg/m<sup>3</sup>, which was too higher than 20 µg/m<sup>3</sup> recommended by WHO guideline. The maximum seasonal mean concentration of PM<sub>10</sub> was observed in winter (138 µg/m<sup>3</sup>). Attributable proportion (AP) due to exposure to PM<sub>10</sub> were estimated 5.487% (95%CI: 3.726-7.185%) of non-accidental mortality cases, 8.011% (95%CI: 4.615-11.174%) of cardiovascular mortality, and 11.174% (95%CI: 4.615-16.215%) of respiratory mortality, for Urmia inhabitants. With every 10 µg/m<sup>3</sup> increase in PM<sub>10</sub> concentrations, relative risk (RR) for total mortality increased by 0.6%.

**Conclusion:** The total cases numbers for hospital admission due to cardiovascular and respiratory diseases in central RR were 257 and 666 cases, respectively. Since incineration fossil fuel, the salty dust storms caused by climate change and drying Urmia Lake is natural, therefore, increasing green space and extensive activities such as desertification are need to control this phenomenon that is extremely challenging.

### Introduction

Population growth, increasing automobiles and development of urbanization have increased atmospheric pollutants worldwide [1]. The global burden of deaths associated with air pollution has been estimated 865,000 annually by World Health Organization (WHO) [2]. Outdoor and

indoor air pollution cause more than 2 million premature deaths each year [3]. Epidemiological studies have demonstrated association between exposure to airborne particulate matter and increased cardiovascular and pulmonary mortality, morbidity from lung cancer, chronic obstructive pulmonary disease (COPD), myocardial infar-

tion and increased respiratory and cardiovascular hospital admissions [4, 5]. International agency for research on cancer (IARC) has defined outdoor air pollution as a mixture and particulate matter specifically as carcinogenic to humans (IARC Group 1). Outdoor air pollution is a combination of different pollutants result from natural and man-made sources [6]. Exposure to particulate matter with the aerodynamic diameter of less than  $10\ \mu\text{m}$  ( $\text{PM}_{10}$ ) cause adverse effects on the human health due to penetration into the human lung. Immune system reactions, lung irritation, bronchitis, asthma exacerbation, hospitalization due to respiratory and cardiovascular diseases, impaired lung function, chronic bronchitis, cancer, and death can be attributed to  $\text{PM}_{10}$  [7]. Small fraction of particulate matters ( $\text{PM}_{2.5}$ , with aerodynamic diameter less than  $2.5\ \mu\text{m}$ ) has more effects on the respiratory tract.  $\text{PM}_{2.5}$  can block the air passages and damage pulmonary mucosal pathways [8]. Increasing every  $10\ \mu\text{g}/\text{m}^3$  in  $\text{PM}_{2.5}$  concentrations in long term exposure has increased 20 and 49% cardiopulmonary mortality and coronary artery deaths, respectively [2]. In addition to size, other physical and chemical characterizations of PM including mass, surface area, number and composition can affect toxicity and health impacts associated with PM [9]. These characterizations are still not conclusively recognized. However, potential health problems may attribute to the organic matters, metals and other chemically active compounds [10]. Recently, drying parts of Lake Urmia, the second saltiest lake in the world, in northwest of Iran (Asian country in the Middle East), has been resulted to releasing a high extent of particulate matter to the air in the West Azerbaijan province and Urmia city as the center of this province. PM,  $\text{SO}_2$  and CO were reported as the most important air pollutants in Urmia city. Dust storm events, inversion, fuel

combustion in vehicles and industries have been introduced as the source of mentioned pollutants [11]. Salty dust related to drying some regions of Urmia Lake releases PM with a significant extent of pollutants such as heavy metals and organic matters and other constituents to the ambient air in the West Azerbaijan province. Combination of PM with heavy metals and different constituents related to Urmia Lake and PM from anthropogenic activities has been anticipated to exacerbate health impacts on population of 3,000,000 people in this province [12-14]. AirQ software is a simple, beneficial and user-friendly tool to assess the effects of criteria air pollutants on the exposed population in a specific place and time. This software has been introduced by the WHO European Centre for Environmental Health, Bilthoven Division [2]. Other researchers have applied AirQ to estimate effect of exposure to different pollutants in Iran [2, 3, 7, 15-19] and other places in the world [20-23]. This study aimed to evaluate exposure to  $\text{PM}_{10}$  related to traffic and drying Lake Urmia and impacts on human health in northwest of Iran according to WHO approach using AirQ. To the best of our knowledge regardless of the significance of salty dust storms on respiratory diseases, this is the first study in term of finding the exposure to  $\text{PM}_{10}$  originating from drying Lake Urmia and impacts on human health in northwest of Iran.

## Materials and methods

### Study area

This is an ecological study which period study was from March 2015 to February 2016. The Urmia city have  $100\ \text{Km}^2$  area and placed in northwest of Iran in longitude of  $45\ 00'$  to  $45\ 07'$  E and latitude of  $37\ 29'$  to  $37\ 34'$  N. This region has arid-cold winters, mild springs, hot dry summers and warm autumn's climate. Sampling sites had four sta-

tions in deferent points if city which they equipped with online air pollutants detector (Enviro SA) by Urmia department of the environment protection (Fig. 1). Also Urmia Lake and wind rose direction in 2015 year illustrate in Fig. 1.

### AirQ 2.2.3

The World health organization software tool AirQ does calculations that allow estimation of the health impact of exposure to air pollution, including assessment of the reduction in life expectancy. AirQ using life-tables method and based on risk assessment from cohort studies evaluations the effects of long-term exposures. This able used to estimate health effects including total mortality, morbidity and hospital admissions. This software related air quality data in various ranges of concentration with epidemiological parameters such as relative risk (RR), baseline incidence (BI in  $10^5$  people), and attributable proportion (AP), and show its results as mortality. The AP in this software is calculated according to the following formula:

$$AP = \text{Sum} \{ [RR(c) - 1] \times p(c) \} / \text{Sum} [RR(c) - 1]$$

Where, RR (c) is the RR of health outcome in the target group, (c) is the population proportion of the target group. By knowing the amount of BI in the target population, we can calculate the attributed value to population contract (IE) by the following formula:

$$IE = BI \times AP$$

Where IE is the quantity of health effect attributable to the exposure, and I is the baseline frequency of the health effect in the target population. Finally, the number of cases attributable to the exposure (NE) can be estimated by the following equation by having the size of population (N) [20].

$$NE = IE \times N$$

RR reflects the rate of a pollutant's impact on the human health by a increas in the pollutant's concentration. RR is found through time-series studies that calculate the concentration increas of air pollutants and their effects on health during a long time period [1].




Fig. 1. Study area, wind rose direction, air pollutants monitoring stations

The concentrations of PM<sub>2.5</sub> and PM<sub>10</sub> from March 2015 to February 2016 were collected from four online air pollution monitoring stations which are situated near to Bahonar, Shaharchi, Azerbaijan, and Motahhari Squares. These monitoring stations are operated by Urmia Department of Environment. The data were filtered by Excel software according to the WHO recommendations. Then, hot and cold seasonal averages, annual averages, and their maximum, as well as annual 98 percentile for PM<sub>2.5</sub> and PM<sub>10</sub>, were extracted. The BI of the total deaths, deaths attributed to cardiovascular and respiratory diseases were estimated using death and diseases data obtained from health Urmia university of medical sciences [24]. In case of needs for other epidemiological parameters which were not available for the city the WHO calculated values were applied. Finally, these data were entered into AirQ software according to the WHO guideline and parameters such as AP, IE, and NE related to PM<sub>10</sub> and PM<sub>2.5</sub> were calculated (Table 1).

## Results and discussion

This research was done AirQ program to estimate the effects of short term exposure to PM<sub>10</sub> concentrations on human health for populations living in urban area. The health effects were assess as the increase in all cases, total mortality, cardiovascular mortality, respiratory mortality, hospital admissions due to cardiovascular diseases, and hospital admissions due to respiratory diseases.

### Concentration of PM<sub>10</sub>

The basic descriptive statistic of PM<sub>10</sub> concentrations were illustrated in Fig. 2. The annual mean of PM<sub>10</sub> concentrations were 106 µg/m<sup>3</sup>, which was too higher than 20 µg/m<sup>3</sup> recommended by WHO as air quality guideline [28]. The maximum concentration value of PM<sub>10</sub> gained during the winter which theirs averages were 138 µg/m<sup>3</sup>, 90 day higher than 20 µg/m<sup>3</sup> and 83 day higher than 50 µg/m<sup>3</sup>. The highest monthly mean of PM<sub>10</sub> concentration with 147 µg/m<sup>3</sup> was recorded in Match. The most probable reasons for this phe-

Table 1. Baseline incidence (BI), relative risk (RR ) with 95% confidence intervals (95% CI), estimated attributable proportion (AP) and number of annual excess cases due to short-term exposure to PM<sub>10</sub> above 10 µg/m<sup>3</sup>

Health Endpoints	pollutant	BI*	RR (per 10 µg/m <sup>3</sup> )	AP percentage (uncertainty range) <sup>b</sup>	Number of excess cases (uncertainty range) <sup>**</sup>
Total mortality	PM <sub>10</sub>	403.82	1.006(1.004-1.008)	5.487(3.726-7.185)	409(278-536)
Cardiovascular Mortality	PM <sub>10</sub>	178.47	1.009(1.005-1.013)	8.011(4.615-11.174)	105(61-146)
Respiratory Mortality	PM <sub>10</sub>	27.62	1.013(1.005-1.020)	11.174(4.615-16.215)	23(10-33)
Hospital Admissions Respiratory Disease	PM <sub>10</sub>	1260	1.008(1.0048-1.0112)	7.185(4.438-9.778)	666(412-907)
Hospital admissions cardiovascular disease	PM <sub>10</sub>	436	1.009(1.006-1.013)	8.011(5.487-11.174)	257(176-359)

\* Crude rate per 100,000 inhabitants per year.

\*\*Obtained using the lower and upper RR values

nomena is windy condition in winter with having the height speed from Lake to city (Fig. 1) and also incineration of fossil fuel with vehicles and residential in the study area. In the other seasons average of  $PM_{10}$  extracted as rank follow:  $101 \mu\text{g}/\text{m}^3$  in autumn  $> 92 \mu\text{g}/\text{m}^3$  in spring  $> 90 \mu\text{g}/\text{m}^3$  in summer. Thus the values of  $PM_{10}$  was rarely in warm seasons less than from could seasons which the reason may be related to inversion potential formation in could seasons and too it lead to intensity of air pollution. The responsible pollutant in many of days in winter reported  $PM$  [29]. Another studies in Yazd, Tehran, Ilam and Isfahan of Iran shown that the cold season having the highest air pollution and responsible pollutant was  $PM$  [8, 10, 15, 19, 30]. The origins of air pollution in these cities mentioned dust storm and traffic which it result similar our finding. Of course in Urmia furthermore dust storm and traffic, drying of Urmia Lake have high potential for salty  $PM$  and air pollution.

### Health effects of $PM_{10}$

The inhabitant's exposure to different ranges of  $PM_{10}$  in the Urmia city was determinate. But, detailed information about confounders were not

easily obtainable. For example, population with sensitive health might be possibly more potential disease to air pollutants, especial particular matters. The days at which populations were exposed to  $PM_{10}$  concentrations were illustrated in Fig. 3. According to this figure the highest values exposure to  $PM_{10}$  taken within range of  $70\text{--}79 \mu\text{g}/\text{m}^3$  during 35 days. Thus, Fig. 3 showed that Urmia population were exposed to  $PM_{10}$  concentrations at a higher than WHO recommended. In Table 1 presented the estimate of AirQ model about the effect of  $PM_{10}$  on human health for total mortality, cardiovascular mortality, respiratory mortality, hospital admissions for cardiovascular diseases, and hospital admissions due to respiratory diseases. Also the annual BI rate of total mortality in Urmia city was 403.82 per 100,000, the cumulative excess cases for total mortality was 105 in intermediate relative risk situations. The number of excess cases predictable for cardiovascular mortality and respiratory mortality was 105 and 23 incidences, respectively. Privies study reported the main mortality reason in Urmia city was belonged to cardiovascular disease with 44.20 percent of total mortality [24].




Fig. 2. Box plot graph of descriptive statistic  $PM_{10}$  concentrations in Urmia, Iran, in 2015 -2016




Fig. 3. Proportion of number cases that are exposed to different concentrations of PM<sub>10</sub>

Fig. 4 show that cases of mortality due to respiratory and cardiovascular diseases in different ranges of PM<sub>10</sub> concentrations which related to 409 cases (RR=1.006) of total mortality, attributable proportion (AP) 5.487% (95%CI: 3.726-7.185%) of non-accidental mortality cases, 8.011% (95%CI; 4.615-11.174%) of cardiovascular mortality, and 11.174% (95%CI; 4.615-16.215%) of respiratory mortality was predictable using AirQ program for the PM<sub>10</sub> concentration more than WHO guideline. With every 10 µg/m<sup>3</sup> rise in PM<sub>10</sub> concentrations, relative risk for total mortality augmented by 0.6%.

On based of Fig. 4, after 59 µg/m<sup>3</sup>, all the mortality was increased very fast. For cardiovascular mortality, with every 10 µg/m<sup>3</sup> increase in PM<sub>10</sub>, RR increased by 0.9%, and for respiratory mortality, 1.3% increase was observed. More than 92% of short-term effects occurred at the time when PM<sub>10</sub> concentration was less than 179 µg/m<sup>3</sup>. In Fig. 6 given the cumulative cases of different health effects attributed to PM<sub>10</sub> concentrations which show the number of hospital admis-

sions (HA) due to cardiovascular diseases and respiratory diseases in different ranges of PM<sub>10</sub> concentration (RR of 1.009 and 1.008, respectively) As this Figure each 10 µg/m<sup>3</sup> increase in PM<sub>10</sub> concentrations has lead 0.8 % increase in RR for HA due to respiratory diseases and cardiovascular diseases.

The total cases numbers for HA due to cardiovascular and respiratory diseases in relative average risk were 257 and 666 persons, respectively. More than 66% of the persons were associated with the PM<sub>10</sub> concentrations of less than 119 µg/m<sup>3</sup>.

In a study in Tehran, exposure to PM<sub>10</sub> concentrations of more than 20 µg/m<sup>3</sup> was reported as major factor for 4.6% of total non-accidental mortality [30]. In another study in Mashad city with 82.9 µg/m<sup>3</sup> value of annual mean PM<sub>10</sub> reported that 4.24% (CI95%; 2.87-5.58) of non-accidental total mortality, 6.23%(CI95%; 3.56-8.7) of cardiovascular mortality, and 8.76% (CI85%; 3.56-12.88) of respiratory mortality are associated to PM<sub>10</sub> concentration [3].




Fig. 4. Estimated cumulative number of total mortality, cardiovascular, and respiratory mortality cases associated PM<sub>10</sub> based on central RR

In other study in urban and industrial area in Tabriz described the total mortality number related to TSP, PM<sub>2.5</sub>, and PM<sub>10</sub> values obtained 327, 360, and 363 µg/m<sup>3</sup>, respectively [31]. In two urban area in Northern Italy reported that PM<sub>2.5</sub> had the main factor for health impact on inhabitants living in two small cities. The exposure to PM<sub>10</sub> in short term with the concentration of more than 20 µg/m<sup>3</sup> in Trieste, a city in the north-east of Italy with about 200,000 population, 52, 28, and 6 cases were assessed for total mortality, car-

diovascular mortality, and respiratory mortality, respectively. They resulted that 2.5% of respiratory mortality and 1.8% of cardiovascular cases were associated to the values of higher than 20 µg/m<sup>3</sup> [32]. The observations of these studies based on excess cases are consistent to that of our study in Urmia. Although further incineration fossil fuel, salty dust from Urmia Lake is sources of PM<sub>10</sub>. Total mortality in Milan from Italy (1,308,000 population), predicted by AirQ model was 677 cases attributable to PM<sub>10</sub> [33].

In totally, a comparison our result with the other researches in Trieste and Tehran could be observed that the high rate of mortality is associated to higher mean values of  $PM_{10}$  or the number of days with high  $PM_{10}$  concentration in the study area due to fuel incineration and also dust from drying Urmia Lake as emerging natural tragedy which it could be treated for population living in some parts of Iran, Turkey, Armenia and Azerbaijan [12]. Furthermore in coming years, it is estimated more than 10 million people will be exposed to this salty PM which may lead to several skin, eye and respiratory diseases.

Also in similar studies in 29 European cities, numerous Asian countries, and 20 American cities, it has been reported that short-term harmful health effects of exposure to  $PM_{10}$  in the developing and developed countries are the same, and mortality rate was heightened by 0.72% when daily  $PM_{10}$  was augmented by  $10 \mu\text{g}/\text{m}^3$  [2, 34-36].

### Conclusion

The findings of this study show that emission pollutants from traffic, incineration of fuel in residential and salty dust from drying of Urmia Lake are main sources for air pollution with  $PM_{10}$ . Thus increasing green space preventive of Lake drying are very emergency.

In this study, the AirQ along with assessment of central point of the relative risk (confidence interval 95%) were applied to consider the health effects of  $PM_{10}$  on Urmia inhabitants in 2015. Results showed that the effect of  $PM_{10}$  on total mortality related to respiratory and cardiovascular disease with 409 cases was more than from many cities in Iran. Also the main mortality reason in Urmia city was belonged to cardiovascular disease with 44.20 percent of total mortality.

Unfortunately the concentration of  $PM_{10}$  was over the concentration than the WHO recommended,

20 in 318 days. Since incineration fossil fuel, the salty dust storms caused by climate change and drying Urmia Lake is natural, therefore, extensive activities such as desertification are needed to control this phenomenon that is extremely challenging. It is clear that such activities are long-term and, therefore, some intermediate and short-term strategies should be carried out to overcome this tragedy.

### Financial supports

Data in this study was taken from Urmia Department of Environment.

### Competing interests

This paper has no competing interests

### Acknowledgements

The authors would like to thank Student Research Committee in Shahid Sadoughi University of Medical Sciences (project number 5527).

### Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

### References

1. Hajizadeh Y, Mokhtari M, Faraji M, Mohammadi A, Nemati S, Ghanbari R, et al. Trends of BTEX in the central urban area of Iran: A preliminary study of photochemical ozone pollution and health risk assessment. *Atmospheric Pollution Research*. 2018;9(2):220-9.
2. Miri M, Aval HE, Ehrampoush MH, Mohammadi A, Toolabi A, Nikonahad A, et al. Human health impact assessment of exposure to particulate matter: an AirQ software modeling. *Environmental Science and Pollution Research*. 2017;24(19):16513-9.
3. Miri M, Derakhshan Z, Allahabadi A, Ahmadi E, Conti GO, Ferrante M, et al. Mortality and morbidity due to exposure to outdoor air pollution in Mashhad metropolis, Iran. The AirQ model approach. *Environmental research*. 2016;151:451-7.



4. Møller P, Danielsen PH, Karotki DG, Jantzen K, Roursgaard M, Klingberg H, et al. Oxidative stress and inflammation generated DNA damage by exposure to air pollution particles. *Mutation Research/Reviews in Mutation Research*. 2014;762:133-66.
5. Künzli N, Kaiser R, Medina S, Studnicka M, Chanel O, Filliger P, et al. Public-health impact of outdoor and traffic-related air pollution: a European assessment. *The Lancet*. 2000;356(9232):795-801.
6. Loomis D, Grosse Y, Lauby-Secretan B, El Ghissassi F, Bouvard V, Benbrahim-Tallaa L, et al. IARC evaluation of the carcinogenicity of outdoor air pollution. *Environnement, Risques & Santé*. 2014;13(4):347-52.
7. Goudarzi G, Daryanoosh S, Godini H, Hopke P, Sicard P, De Marco A, et al. Health risk assessment of exposure to the Middle-Eastern Dust storms in the Iranian megacity of Kermanshah. *Public Health*. 2017;148:109-16.
8. Mokhtari M, Miri M, Mohammadi A, Khorsandi H, Hajizadeh Y, Abdolhnejad A. Assessment of air quality index and health impact of PM<sub>10</sub>, PM<sub>2.5</sub> and SO<sub>2</sub> in Yazd, Iran. *Journal of Mazandaran University of Medical Sciences*. 2015;25(131):14-23.
9. Osornio-Vargas AR, Serrano J, Rojas-Bracho L, Miranda J, García-Cuellar C, Reyna MA, et al. In vitro biological effects of airborne PM<sub>2.5</sub> and PM<sub>10</sub> from a semi-desert city on the Mexico-US border. *Chemosphere*. 2011;83(4):618-26.
10. Naddafi K, Atafar Z, Faraji M, Ghanbarian M, Rezaei S, Ghoskhal MG, et al. Health effects of airborne particulate matters (PM<sub>10</sub>) during dust storm and non-dust storm conditions in Tehran. *Journal of Air Pollution and Health*. 2016;1(4):259-68.
11. Khorsandi H, Amini Tapok F, Cargar H, Mousavi Moughanjogi S. Study of Urmia city air quality according to the air quality index (AQI). *Urmia Medical Journal*. 2013;23(7):767-75.
12. Hajizadeh Y, Namati S, Aghapour AA, Abdolhnejad A, Mohammadi A, Panahi H, et al. Influence of Air Pollution on Chemical Quality of Wet Atmospheric Deposition: a Case Study in Urmia, Iran. *Iranian Journal of Health, Safety and Environment*. 2017;5(1):904-10.
13. Gholampour A, Nabizadeh R, Hassanvand MS, Taghipour H, Nazmara S, Mahvi AH. Characterization of saline dust emission resulted from Urmia Lake drying. *Journal of Environmental Health Science and Engineering*. 2015;13(1):82.
14. Gholampour A, Nabizadeh R, Hassanvand MS, Nazmara S, Mahvi AH. Elemental composition of particulate matters around Urmia Lake, Iran. *Toxicological & Environmental Chemistry*. 2017;99(1):17-31.
15. Abdolhnejad A, Jafari N, Mohammadi A, Miri M, Hajizadeh Y, Nikoonahad A. Cardiovascular, respiratory, and total mortality ascribed to PM<sub>10</sub> and PM<sub>2.5</sub> exposure in Isfahan, Iran. *Journal of Education and Health Promotion*. 2017;6(1):109.
16. Ghoskhal MG, Heibati B, Naddafi K, Kloog I, Conti GO, Polosa R, et al. Evaluation of chronic obstructive pulmonary disease (COPD) attributed to atmospheric O<sub>3</sub>, NO<sub>2</sub>, and SO<sub>2</sub> using Air Q Model (2011–2012 year). *Environmental research*. 2016;144:99-105.
17. Khaniabadi YO, Daryanoosh SM, Hopke PK, Ferrante M, De Marco A, Sicard P, et al. Acute myocardial infarction and COPD attributed to ambient SO<sub>2</sub> in Iran. *Environmental Research*. 2017;156:683-7.
18. Khaniabadi YO, Goudarzi G, Daryanoosh SM, Borgini A, Tittarelli A, De Marco A. Exposure to PM<sub>10</sub>, NO<sub>2</sub>, and O<sub>3</sub> and impacts on human health. *Environmental Science and Pollution Research*. 2017;24(3):2781-9.
19. Nikoonahad A, Naserifar R, Alipour V, Poursafar A, Miri M, Ghafari HR, et al. Assessment of hospitalization and mortality from exposure to PM10 using AirQ modeling in Ilam, Iran. *Environmental Science and Pollution Research*. 2017;24(27):21791-6.
20. Fattore E, Paiano V, Borgini A, Tittarelli A, Bertoldi M, Crosignani P, et al. Human health risk in relation to air quality in two municipalities in an industrialized area of Northern Italy. *Environmental research*. 2011;111(8):1321-7.
21. Jeong SJ. The impact of air pollution on human health in Suwon City. *Asian Journal of Atmospheric Environment*. 2013;7(4):227-33.
22. Kim S-Y, Lee J-T, Hong Y-C, Ahn K-J, Kim H. Determining the threshold effect of ozone on daily mortality: an analysis of ozone and mortality in Seoul, Korea, 1995–1999. *Environmental research*. 2004;94(2):113-9.
23. Lee YJ, Lim YW, Yang JY, Kim CS, Shin YC, Shin DC. Evaluating the PM damage cost due to urban air pollution and vehicle emissions in Seoul, Korea. *Journal of environmental management*. 2011;92(3):603-9.
24. Majidi M, Ghaemi S, Javaezi M, Delirrad M. A Study on Human Mortalities & Death Registration System in West Azerbaijan Province of Iran. *IJFM. Iranian journal of forensic medicine*. 2016;22(3):193-202.
25. Urmia municipality. <http://eng.urmia.ir/Default.aspx>.
26. Miri M, Shendi MRA, Ghaffari HR, Aval HE, Ahmadi E, Taban E, et al. Investigation of outdoor BTEX: Concentration, variations, sources, spatial distribution, and risk assessment. *Chemosphere*. 2016;163:601-9.
27. kurdpress News. <http://www.kurdpress.com/details.aspx?id=5800>.
28. WHO. Air Quality Guidelines: Global Update 2005. Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide: World Health Organization; 2006.
29. Department of Environmental Protection, Islamic Republic of Iran. [Internet]. 2017. Available from: <https://www.doe.ir/Portal/home>.
30. Naddafi K, Hassanvand MS, Yunesian M, Momeniha F, Nabizadeh R, Faridi S, et al. Health impact assessment of air pollution in megacity of Tehran, Iran. *Iranian journal of environmental health science & engineering*. 2012;9(1):28.
31. Gholampour A, Nabizadeh R, Naseri S, Yunesian M, Taghipour H, Rastkari N, et al. Exposure and health impacts of outdoor particulate matter in two urban and

- industrialized area of Tabriz, Iran. *Journal of Environmental Health Science and Engineering*. 2014;12(1):27.
32. Tominz R, Mazzoleni B, Daris F. Estimate of potential health benefits of the reduction of air pollution with PM<sub>10</sub> in Trieste, Italy. *Epidemiologia e prevenzione*. 2005;29(3-4):149-55.
33. Martuzzi M, Galassi C, Forastiere F, Bertollini R. Health impact assessment of air pollution in the eight major Italian cities. Health impact assessment of air pollution in the eight major Italian cities. OMS. 2002.
34. Committee HIO. Health effects of outdoor air pollution in developing countries of Asia: a literature review. Health Effects Institute, Boston, USA. 2004.
35. Ezzati M. Comparative quantification of health risks: sexual and reproductive health: World Health Organization. 2004;2:1177-320.
36. Samet JM, Zeger SL, Dominici F, Curriero F, Coursac I, Dockery DW, et al. The national morbidity, mortality, and air pollution study. Part II: morbidity and mortality from air pollution in the United States *Res Rep Health Eff Inst*. 2000;94(pt 2):5-79.