



Seasonal trend analysis and influence of meteorological factors on the concentration of atmospheric pollutants in an urban area

Naresh Kumar¹, Priya Ranjan², Suman^{2,*}

¹ Atmospheric Research Laboratory, Department of Environmental Sciences, Sharda School of Basic Sciences and Research, Sharda University, Greater Noida, India

² Department of Environmental Sciences, Sharda School of Basic Sciences and Research, Greater Noida, India

ARTICLE INFORMATION

Article Chronology:

Received 05 April 2024

Revised 10 September 2024

Accepted 12 December 2024

Published 30 December 2024

Keywords:

Meteorological variable; Pearson correlation; Air pollutants; Linear relation; National ambient air quality standards (NAAQS)

CORRESPONDING AUTHOR:

suman.ism3@gmail.com

Tel: (+91 120) 4570000

Fax: (+91 120) 4570000

ABSTRACT

Introduction: Air pollution is a growing concern in Greater Noida, Uttar Pradesh, where meteorological factors play a major role. As urbanization accelerates, the region has not yet thoroughly examined the impact of factors like temperature, humidity, wind direction, and speed on pollutants like Particulate Matters (PM_{2.5}, PM₁₀), Nitrogen dioxide (NO₂), and Sulfur dioxide (SO₂)

Materials and methods: To explore this issue, a 12-month study (January to December 2023) was conducted to analyze the relationships between these meteorological factors and pollutant levels. In the present study, pollutant movement was tracked over time. Pearson correlation was used to correlate air pollution concentrations with weather conditions.

Results: The study revealed that the average concentrations of PM₁₀, PM_{2.5}, and NO₂ were 912.54 µg/m³, 257.42 µg/m³, and 65.98 µg/m³, respectively. In contrast, SO₂ had an average concentration of 26.85 µg/m³.

Conclusion: The study indicates that particulate matter and temperature are strongly negatively correlated, while wind speed shows a weak positive correlation with particulate matter. Pollution levels and relative humidity also display a negative correlation, as do particulate matter concentrations and humidity. However, gaseous pollutants exhibit little to no significant correlation with these weather conditions

Please cite this article as: Kumar N, Ranjan P, Suman. Seasonal trend analysis and influence of meteorological factors on the concentration of atmospheric pollutants in an urban area. Journal of Air Pollution and Health. 2024;9(4): 493-508.



Introduction

The spatial variation, diffusion, and transport of pollutants are largely influenced by the geographical distribution of emission sources, with meteorological factors playing a crucial role. The National Air Quality Monitoring Program (NAMP) highlights that various climatic factors play a role in the dispersion and persistence of air pollutants. When atmospheric emissions surpass certain thresholds, they become pollutants, a process that can occur due to higher emissions and specific meteorological situations. Unlike emissions, which can be regulated, weather conditions remain beyond our control [1, 2]. For example, research in New Delhi, India, linked severe air pollution events to stagnant weather patterns [3].

It has been identified atmospheric impairment as one of the major ecological issues globally. It is also a rising issue in low-economic countries in South Asia, such as India, which are rapidly industrializing [3]. Fine Particulate Matter (FPM) has been recognized as the most serious environmental hazard due to its life-threatening toxicity to people's health. Municipal air contamination is a serious and growing ecological problem, particularly in emerging countries, and it significantly impacts public health worldwide, specifically respiratory and cardiac diseases.

The early mortality due to atmospheric impairment is the leading cause of death globally. In India, air contamination causes approximately 16.7 lakh untimely deaths, accounting for around eighteen percent of total mortalities. Eleven out of the twelve most polluted cities in the world with extreme PM concentrations are in India, raising serious concerns [4].

The quality of air in cities is affected by several factors such as micrometeorological conditions and urban growth trends. The combined impact of these factors can be more harmful to one's health than the impact of each factor on its own. The increasing occurrence of life-threatening

weather events due to climate change caused by human activities is a cause for concern. To mitigate the negative effects of these factors, it is crucial to understand the relationships between different areas within a city, verify weather patterns, predict air quality, and control contamination [5]. Meteorological conditions also contribute to the increase of air impurities instead of just emissions into the atmosphere. Various studies have shown that factors such as temperature, humidity, air flow direction, and velocity have an effect on pollution concentrations within local areas.

Climatology and topography are the two major aspects that influence the occurrence of contamination [6], the extent of exposure, and the effects of pollution [7]. The dispersion, residence time, chemical transformations, and transport of contaminants in the air are all influenced by meteorology [8]. The levels of heavy metal-containing particulate matter are influenced by climatological factors such as speed and direction of the wind, temperature, and dampness [9]. The most severe exposure episodes occur predominantly during prolonged areas of low wind speeds and atmospheric stability, which favor secondary aerosol generation near the source locations [10]. According to numerous research, the largest amounts of particulate matter are seen in winter, followed by summer and the rainy season [11].

In some cases, daily levels of air pollutants are strongly influenced by weather factors such as wind speed and solar radiation [12], while in other cases, humidity and cloud cover can have an impact, particularly during the summer months for pollutants like Sulfur dioxide (SO₂) [13]. Nitrogen dioxide (NO₂) levels are strongly linked with temperature during the monsoon season and humidity and mist cover during winter.

Research has generally indicated a weak but significant relationship between PM₁₀ levels and weather conditions. In one study conducted in Coimbatore, India, six different pollutants were

examined in manufacturing, housing, rush-hour traffic, and commercial areas. Although tropospheric ozone and carbon monoxide showed weak correlations with temperature and wind speed, they did show a positive correlation with temperature and a negative correlation with humidity [14].

Meteorological elements like temperature, humidity, wind speed, and rainfall play a crucial role in influencing atmospheric pollutants. Understanding and monitoring these factors are essential for controlling pollution levels in urban settings. Various studies have shown the impact of weather conditions on air quality. However, while these studies have provided insights into the general air quality in Greater Noida, they have not specifically analyzed key pollutants across different seasons or explored the seasonal variations in detail [15-18].

This study aims to bridge the gap by offering a thorough analysis of air quality trends in Greater Noida from January to December 2023. It seeks to pinpoint significant changes and variations, providing insights into the success of pollution control measures and identifying any new pollution sources. Additionally, the study emphasizes the crucial role of meteorological factors in influencing air quality, highlighting the complex interactions between weather conditions, pollutant dispersion, and concentrations. The primary goal is to examine the relationship between pollutant levels and meteorological conditions and to evaluate the seasonal variations in air pollution in Greater Noida.

Materials and methods

Study area and sampling sites

Greater Noida, situated in the Gautam Buddha Nagar district of Uttar Pradesh, India, is positioned at 28.474400° N and 77.504000° E, spanning an area of around 40,000 hectares. As of March 2014, the city had a population of 107,676 across 124 localities and is part of the Delhi National Capital Region (NCR).

The climate in Greater Noida is similar to Delhi, featuring hot and dry summers, wet monsoons, and pleasant spring and autumn seasons. Classified as Seismic Zone III by the Bureau of Indian Standards, the city has a moderate earthquake risk. Additionally, a United Nations Development Programme (UNDP) assessment indicates its vulnerability to severe wind and cyclone damage.

Greater Noida experiences a tropical savanna climate with three main seasons: summer, monsoon, and winter. During summer (March to June), temperatures range from 23°C to 45°C. The monsoon season, from mid-June to mid-September, brings an average rainfall of 93.2 cm (36.7 in). In winter, temperatures can drop to 3–4°C, and dense fog often reduces visibility in January. The area was selected for the study due to its strategic importance as a growing hub for trade, tourism, population growth, and communication.

However, increasing traffic has led to a decline in air quality, contributing to rising pollution levels. The study was conducted at eight different locations in Greater Noida to measure concentrations of various air pollutants, as detailed in Table 1 and illustrated in Fig. 1.

Table 1. Sampling sites in Greater Noida

Site No.	Study Site	Category	Location	
S1	Pari Chowk	Major Road	28°27'46.07" N	77°31'00.31" E
S2	Banghel	Residential Road	28°36'32.76" N	77°26'44.93" E
S3	Sharda University	Arbitrary Road	28°28'25.14" N	77°29'03.89" E
S4	National Highway-91	Major Road	26°33'41.47" N	80°14'18.81" E
S5	Greater Noida West Link Road	Major Road	28°29'51.61" N	77°32'07.04" E
S6	Dadri Main Road	Major Road	28°32'10.66" N	77°25'19.12" E
S7	Alpha 1 Commercial Complex	Commercial Road	28°28'16.29" N	77°30'43.05" E
S 8	Amit Nagar, Sadarpur	Residential Road	28°27'52.30" N	77°32'28.21" E




Fig. 1. Location map of study area

Table 2. Methods and instruments used for ambient air quality assessment

Parameter	Instrument used	Measurement method
PM ₁₀	Respirable Dust Sampler (APM 460 NL)	Gravimetric Methodology IS 5182(Part 23): 2006
PM _{2.5}	Fine Particulate Sampler (APM 550 MFC)	Gravimetric Methodology IS 5182(Part 24): 2019
NO ₂	Gaseous Pollutant Sampler (APM 433)	Modified Jacobs and Hochheiser method IS 5182(Part 6): 2017
SO ₂	Gaseous Pollutant Sampler (APM 433)	Modified West & Gaeke Method IS 5182 (Part 2): 2001

Instrumentation and collection of data

The standard protocols were used for the ambient particulate and gaseous pollutants which are detailed in Table 2.

The primary objective was focused on the relationship between air pollution and weather in Greater Noida. Data on temperature, humidity, wind direction, and airspeed were gathered from Central Pollution Control Board (CPCB)'s online portal. CPCB ensures high data quality through rigorous sampling, review, and standardization processes.

Statistical analysis

The study analyzed the spatial distribution of air pollutants using statistical methods. Data processing was done with SPSS, while mapping and visualization were carried out using Origin Pro and SPSS. The Pearson correlation coefficient was employed to evaluate the relationship between weather factors and pollutant levels, indicating the strength and direction of their linear relationship, with values ranging from -1 to +1. A value of 0 signifies no correlation [19].

Results and discussion

The study examined particulate matter and

gaseous pollutants at eight monitoring sites in Greater Noida. Results are shown in Table 3. The study period was divided into three seasons: winter (January-March), summer (April-June), and post-monsoon (October-December).

Fig. 2 shows seasonal variations in particulate matter at the eight sites. In winter, Dadri Main Road had the highest PM₁₀ concentration at 912.54 µg/m³, while Pari Chowk had the lowest at 240.47 µg/m³. The highest PM_{2.5} levels were also recorded in winter due to lower mixing layer heights, which cause pollutants to accumulate near the ground. This accumulation is driven by high emissions from vehicles, manufacturing, and fossil fuel combustion. The Mixing Layer Height (MLH) is a key factor that influences pollutant concentration near the surface.

The Dadri Main Road (912.54 µg/m³) had the greatest mean concentration of PM₁₀, followed by Bhangel (684.96 µg/m³) and Amit Nagar (478.57 µg/m³), while the Pari Chowk, Alpha 1 commercial complex, and Sharda University had the lowest. (340.45 µg/m³). Similar to PM_{2.5}, the greatest average concentration was found at Dadri Main Road (257.42 µg/m³), followed by Bhangel (141.13 µg/m³), and Sharda University (103.66 µg/m³), while the lowest concentration was found at Amit Nagar, Sadarpur. (35.63 µg/m³).

Table 3. Seasonal changes in air pollutants in the study area

Site	Pollutants (in ug/m ³)	Seasons		
		Winter	Summer	Post- Monsoon
S1	PM ₁₀	240.47	336.72	254.68
	PM _{2.5}	93.99	73.49	104.61
	NO ₂	15.62	13.40	17.24
	SO ₂	16.36	34.49	30.57
S2:	PM ₁₀	684.96	585.71	650.78
	PM _{2.5}	141.13	78.74	135.94
	NO ₂	40.54	15.69	26.46
	SO ₂	24.55	35.64	28.93
S3:	PM ₁₀	340.45	406.46	385.38
	PM _{2.5}	103.66	68.33	86.24
	NO ₂	20.42	21.75	24.56
	SO ₂	16.76	30.61	27.36

Table 3. Continued

Site	Pollutants (in $\mu\text{g}/\text{m}^3$)	Seasons		
		Winter	Summer	Post- Monsoon
S4:	PM ₁₀	372.45	472.16	514.23
	PM _{2.5}	88.30	75.56	105.36
	NO ₂	22.59	20.72	28.56
	SO ₂	26.58	32.65	26.86
S5:	PM ₁₀	334.43	380.62	426.35
	PM _{2.5}	40.47	66.65	110.35
	NO ₂	62.27	10.49	54.38
	SO ₂	19.53	34.63	40.38
S6:	PM ₁₀	912.54	536.51	754.23
	PM _{2.5}	257.42	73.37	153.85
	NO ₂	12.55	10.69	15.68
	SO ₂	10.89	32.65	34.59
S7:	PM ₁₀	299.48	518.20	350.57
	PM _{2.5}	92.40	78.23	105.68
	NO ₂	22.53	10.62	28.35
	SO ₂	18.53	34.63	36.45
S8:	PM ₁₀	478.57	250.68	385.13
	PM _{2.5}	35.63	36.54	75.68
	NO ₂	24.71	10.50	31.86
	SO ₂	13.97	43.66	34.58




Fig. 2. Seasonal variation of ambient particulate matter in study area




Fig. 2. Continued

Variability of ambient gaseous pollutants

The data in Fig. 3 highlights that the Noida-Greater Noida highway experienced the highest nitrogen dioxide levels in the winter, with Greater Noida following closely in the post-monsoon season. In comparison, the lowest nitrogen dioxide concentration was observed during the summer in Amit Nagar, Sadarpur. The primary reason for the higher concentration during the winter season is that the dispersion of gaseous pollutants is inversely proportional to temperature and wind speed, which means that cooler temperatures and lower wind speeds lead to poor dispersion of pollutants.

Variability of ambient gaseous pollutants

The data in Fig. 3 highlights that the Noida-Greater Noida highway experienced the highest nitrogen dioxide levels in the winter, with Greater Noida following closely in the post-monsoon season. In comparison, the lowest nitrogen dioxide

concentration was observed during the summer in Amit Nagar, Sadarpur. The primary reason for the higher concentration during the winter season is that the dispersion of gaseous pollutants is inversely proportional to temperature and wind speed, which means that cooler temperatures and lower wind speeds lead to poor dispersion of pollutants.

According to the meteorological data collected from the CPCB portal, the wind speed in the study region varies from 0.00 to 5.43 m/s with an average speed of 1.46 m/s throughout the study tenure. During the study episode mainly North-Easterly, North-North Easterly, and East-North Easterly winds prevailed in the study area. To represent the wind flow in the study area a wind rose diagram was plotted by using WRPLOT View – Lakes Environmental Software which is shown in Fig. 5. The wind rose plot is imposed on satellite imagery of the study area by using Google Earth which is shown in Fig. 6 to show the flowing direction of the wind in the study area during different seasons.




Fig. 3. Seasonal variation of gaseous pollutants in study area

Table 4. Variation of meteorological parameters in the study area

		Range	Minimum	Maximum	Mean	Standard Deviation	Variance	Skewness	Kurtosis
Winter	Temperature	8.51	26.92	35.43	31.60	1.57	2.46	-1.36	1.97
	Relative Humidity	36.40	43.48	79.88	63.46	8.40	70.54	0.039	-0.646
	Wind Speed	3.83	0.30	4.13	1.07	0.73	0.5	2.0	4.36
	Wind Direction	121.53	38.50	160.03	83.88	31.01	961.80	0.82	-0.356
Summer	Temperature	10.07	27.93	38.00	32.10	2.45	6.01	0.130	-1.045
	Relative Humidity	40.00	26.00	66.00	39.70	10.32	106.58	0.900	-0.098
	Wind Speed	5.43	0.00	5.43	2.41	1.05	1.11	0.44	0.04
	Wind Direction	103.83	47.17	151.00	95.16	29.15	849.91	0.25	-1.18
Post-Monsoon	Temperature	6.00	29.00	35.00	31.37	1.22	1.49	0.58	1.06
	Relative Humidity	33.00	49.00	82.00	62.23	8.12	65.91	0.65	0.24
	Wind Speed	2.00	0.00	2.00	0.90	0.51	0.26	-0.17	0.85
	Wind Direction	261.00	45.00	306.00	102.78	54.11	2927.87	2.24	5.27




Fig. 4. Variation of meteorological parameters in the study area




Fig. 5. Wind rose diagram for different seasons in the study area

Relationship between air pollutants and meteorological factors

The correlation analysis was carried out to explore the relationships between ambient air pollutant levels and meteorological factors such as temperature, humidity, wind speed, and wind direction, as illustrated in Table 5. The findings revealed that PM₁₀ had the weakest negative correlation with all meteorological parameters, with correlation coefficients of -0.194 for temperature, -0.180 for humidity, -0.154 for wind speed, and -0.422 for wind direction. Higher temperatures contribute to increased mixing heights, resulting from greater heat flux and thermal turbulence, which helps disperse pollutants and reduce their concentrations in the air.

A consistent negative correlation between pollutant levels and relative humidity was observed throughout the study. Both PM_{2.5} and PM₁₀ concentrations were inversely correlated with humidity, with correlation values of $r = -0.180$ and $r = -0.413$, respectively, while gaseous pollutants showed strong positive correlations ($r = 0.881$ and $r = 0.825$). The negative relationship between PM₁₀ and humidity can be attributed to particle clumping, as increased humidity causes particles to combine, grow larger, and settle out of the atmosphere through dry deposition. However, this pattern did not extend to gaseous pollutants.

Similar studies support this negative correlation between particulate matter and humidity [25].

PM₁₀ also demonstrated a negative correlation with wind speed ($r = -0.154$), while PM_{2.5} had a slight positive correlation ($r = 0.228$). Gaseous pollutants, however, exhibited weak or non-significant negative correlations with wind speed, with $r = -0.080$ and $r = -0.350$, respectively. Low wind speeds were linked to higher pollutant concentrations due to atmospheric stability, which hinders the dispersion of pollutants [26]. Wind speed had minimal positive correlations with particulate matter, but for PM_{2.5}, the relationship was moderate ($r = 0.270$). Gaseous pollutants displayed weak negative correlations with airspeed ($r = -0.219$ and $r = -0.145$).

The inverse relationships between wind speed, humidity, and particulate matter suggest that both wind and humidity aid in dispersing pollutants. However, high wind speeds might also resuspend dust particles, leading to increased PM levels. Additionally, the positive correlation between wind direction and particulate matter indicates that ground-level dust resuspension likely contributed to elevated PM concentrations at the study site. Sunlight and temperature were also found to influence gaseous pollutants, with lower temperatures favoring particulate matter deposition in the atmosphere [27].

Table 5. Correlation between ambient air pollutants and meteorological parameters

		PM ₁₀	PM _{2.5}	NO ₂	SO ₂	Temperature	Relative Humidity	Wind Speed	Wind Direction
PM ₁₀	Pearson Correlation	1	.840**	-.183	-.176	-.194	-.180	-.154	-.422
	Sig. (2-tailed)		.009	.664	.678	.645	.669	.716	.297
	N	8	8	8	8	8	8	8	8
PM _{2.5}	Pearson Correlation	.840**	1	-.445	-.502	-.026	-.413	.228	.093
	Sig. (2-tailed)	.009		.270	.205	.951	.309	.587	.826
	N	8	8	8	8	8	8	8	8
NO ₂	Pearson Correlation	-.183	-.445	1	.660	.153	.881**	-.080	-.540
	Sig. (2-tailed)	.664	.270		.075	.717	.004	.850	.167
	N	8	8	8	8	8	8	8	8
SO ₂	Pearson Correlation	-.176	-.502	.660	1	-.259	.825*	-.350	-.447
	Sig. (2-tailed)	.678	.205	.075		.536	.012	.396	.267
	N	8	8	8	8	8	8	8	8

Table 5. Continued

		PM ₁₀	PM _{2.5}	NO ₂	SO ₂	Temperature	Relative Humidity	Wind Speed	Wind Direction
Temperature	Pearson Correlation	-.194	-.026	.153	-.259	1	.042	.681	.333
	Sig. (2-tailed)	.645	.951	.717	.536		.922	.063	.420
	N	8	8	8	8	8	8	8	8
Relative Humidity	Pearson Correlation	-.180	-.413	.881**	.825*	.042	1	-.141	-.413
	Sig. (2-tailed)	.669	.309	.004	.012	.922		.739	.310
	N	8	8	8	8	8	8	8	8
Wind Speed	Pearson Correlation	-.154	.228	-.080	-.350	.681	-.141	1	.524
	Sig. (2-tailed)	.716	.587	.850	.396	.063	.739		.183
	N	8	8	8	8	8	8	8	8
Wind Direction	Pearson Correlation	-.422	.093	-.540	-.447	.333	-.413	.524	1
	Sig. (2-tailed)	.297	.826	.167	.267	.420	.310	.183	
	N	8	8	8	8	8	8	8	8

** Correlation is significant at the 0.01 level (2-tailed)

*Correlation is significant at the 0.05 level (2-tailed)




Fig. 6. Correlation matrix between ambient air pollutants and meteorological parameters

Conclusion

Throughout the study period, PM₁₀, PM_{2.5}, and NO₂ levels consistently surpassed the annual standards set by the Central Pollution Control Board (CPCB). Additionally, the 24-hour concentrations of these pollutants exceeded permissible limits, indicating deteriorating air quality in Greater Noida. Conversely, SO₂ levels remained within both the annual and 24-h limits set by the CPCB, likely due to effective government policies such as promoting LPG over coal for household fuel, reducing sulfur content in diesel, converting diesel vehicles to CNG, and enforcing Bharat Stage-VI emission standards.

Seasonal trends were observed for PM_{2.5}, PM₁₀, SO₂, and NO₂, with the highest concentrations during winter and the lowest in summer. This variation can be partially attributed to seasonal changes in the atmospheric boundary layer, which is lower in winter and higher in summer. These findings highlight the significant impact of meteorological factors on air pollutant concentrations.

Pearson correlation analysis revealed significant relationships between pollutants. PM₁₀ and PM_{2.5} showed a strong correlation, suggesting common sources. Notable correlations were also found between air pollutants and meteorological variables. Wind speed was inversely related to pollutant levels, emphasizing the role of wind in dispersing pollutants. Similarly, temperature was inversely correlated with pollutant concentrations, indicating that increased heat flux and thermal turbulence help reduce pollutant levels in the atmosphere.

This study provides valuable insights into how meteorological conditions influence air pollution, offering a foundation for

policymakers to develop effective pollution control strategies. It also underscores the need to consider the impact of global climate change on air quality and public health. While the study focused on surface-level meteorological factors, it suggests that the vertical structure of the atmospheric boundary layer may also play a crucial role in influencing ground-level pollutant concentrations.

Financial supports

This research does not receive any funding from external sources.

Competing interests

The authors declare no conflict of interest.

Authors contributions

Suman (corresponding author): Oversight, conceptual development, methodology design, investigation, visualization, project management, validation, drafting the original manuscript, and reviewing and editing. Naresh Kumar (lead author): Conceptualization, methodology, investigation, and drafting of the original manuscript. Priya Ranjan: Involved in conceptualization, methodology, and investigation.

Acknowledgements

The researchers extend their heartfelt thanks to Sharda University, Greater Noida, for offering the essential laboratory facilities required for this study. They also express their gratitude to the anonymous reviewers for their insightful feedback and expertise, which greatly improved the quality of this manuscript.

Ethical considerations

The researchers have strictly followed all ethical guidelines, covering aspects such as plagiarism, informed consent, research misconduct, data fabrication or falsification, duplicate publication or submission, and redundancy.

References

1. Cheng N, Cheng B, Li S, Ning T. Effects of meteorology and emission reduction measures on air pollution in Beijing during heating seasons. *Atmospheric Pollution Research*. 2019 May 1;10(3):971-9.
2. Ma T, Duan F, He K, Qin Y, Tong D, Geng G, et al. Air pollution characteristics and their relationship with emissions and meteorology in the Yangtze River Delta region during 2014–2016. *Journal of Environmental Sciences*. 2019 Sep 1;83:8-20.
3. Kanawade VP, Srivastava AK, Ram K, Asmi E, Vakkari V, Soni VK, et al. What caused severe air pollution episode of November 2016 in New Delhi?. *Atmospheric Environment*. 2020 Feb 1;222:117125.
4. Shi Y, Bilal M, Ho HC, Omar A. Urbanization and regional air pollution across South Asian developing countries—A nationwide land use regression for ambient PM_{2.5} assessment in Pakistan. *Environmental Pollution*. 2020 Nov 1;266:115145.
5. Ravindra K, Singh T, Mor S, Singh V, Mandal TK, Bhatti MS, et al. Real-time monitoring of air pollutants in seven cities of North India during crop residue burning and their relationship with meteorology and transboundary movement of air. *Science of the total environment*. 2019 Nov 10;690:717-29.
6. Ravindra K, Singh T, Sinha V, Sinha B, Paul S, Attri SD, et al. Appraisal of regional haze event and its relationship with PM_{2.5} concentration, crop residue burning and meteorology in Chandigarh, India. *Chemosphere*. 2021 Jun 1;273:128562.
7. Singh T, Biswal A, Mor S, Ravindra K, Singh V, Mor S. A high-resolution emission inventory of air pollutants from primary crop residue burning over Northern India based on VIIRS thermal anomalies. *Environmental Pollution*. 2020 Nov 1;266:115132.
8. Ulpiani G, Hart MA, Di Virgilio G, Maharaj AM. Urban meteorology and air quality in a rapidly growing city: Inter-parameter associations and intra-urban heterogeneity. *Sustainable Cities and Society*. 2022 Feb 1;77:103553.
9. Hertig E, Russo A, Trigo R. Compound heat and ozone pollution events relevant for human health. In: *EGU General Assembly Conference Abstracts 2021 Apr* (pp. EGU21-119).
10. O'Lenick CR, Wilhelmi OV, Michael R, Hayden MH, Baniassadi A, Wiedinmyer C, et al. Urban heat and air pollution: A framework for integrating population vulnerability and indoor exposure in health risk analyses. *Science of the total environment*. 2019 Apr 10;660:715-23.
11. Pandey A, Brauer M, Cropper ML, Balakrishnan K, Mathur P, Dey S, et al. Health and economic impact of air pollution in the states of India: the Global Burden of Disease Study 2019. *The Lancet Planetary Health*. 2021 Jan 1;5(1):e25-38.
12. Tandon A, Kanchan T, Tandon A. The poison we breathe. *The Lancet*. 2020 Feb 1;395(10221):e18.

13. Khaiwal Ravindra KR, Neha Agarwal NA, Kaur-Sidhu M, Suman Mor SM. Appraisal of thermal comfort in rural household kitchens of Punjab, India and adaptation strategies for better health.
14. Sigamani S, Venkatesan R. Air quality index prediction with influence of meteorological parameters using machine learning model for IoT application. *Arabian Journal of Geosciences*. 2022 Feb;15(4):340.
15. Briceno-Garmendia C, Qiao W, Foster V. The economics of electric vehicles for passenger transportation. World Bank Publications; 2023 Jun 6.
16. Bose A, Roy Chowdhury I. Investigating the association between air pollutants' concentration and meteorological parameters in a rapidly growing urban center of West Bengal, India: a statistical modeling-based approach. *Modeling Earth Systems and Environment*. 2023 Jun;9(2):2877-92.
17. Cai X, Yu J, Qin Y. Spatial Distribution of Air Pollution and Its Relationship with Meteorological Factors: A Case Study of 31 Provincial Capitals in China. *Polish Journal of Environmental Studies*. 2023 May 1;32(3).
18. Banerjee B, Kundu S, Kanchan R, Mohanta A. Examining the relationship between atmospheric pollutants and meteorological factors in Asansol city, West Bengal, India, using statistical modelling. *Environmental Science and Pollution Research*. 2024 May 18:1-8.
19. Singh D, Kumar AS, Goyal VC, Arora M, Allaka NR. Characteristics of meteorological variables and their implications on evaporation in Roorkee (India). *HydroResearch*. 2021 Jan 1;4:47-60.
20. Javanroodi K, Nik VM. Interactions between extreme climate and urban morphology: Investigating the evolution of extreme wind speeds from mesoscale to microscale. *Urban Climate*. 2020 Mar 1;31:100544.
21. Oji S, Adamu H. Correlation between air pollutants concentration and meteorological factors on seasonal air quality variation. *Journal of air pollution and health*. 2020 May 27.
22. Feng H, Zou B, Wang J, Gu X. Dominant variables of global air pollution-climate interaction: Geographic insight. *Ecological indicators*. 2019 Apr 1;99:251-60.
23. Battista G, de Lieto Vollaro R. Correlation between air pollution and weather data in urban areas: Assessment of the city of Rome (Italy) as spatially and temporally independent regarding pollutants. *Atmospheric Environment*. 2017 Sep 1;165:240-7.
24. Hrishikesh CG, Nagendra SS. Study of meteorological impact on air quality in a humid tropical urban area. *Journal of Earth System Science*. 2019 Jul;128:1-8.
25. Manju A, Kalaiselvi K, Dhananjayan V, Palanivel M, Banupriya GS, Vidhya MH, et al. Spatio-seasonal variation in ambient air pollutants and influence of meteorological factors in Coimbatore, Southern India. *Air Quality, Atmosphere and Health*. 2018 Dec;11:1179-89.
26. Seinfeld JH, Pandis SN. *Atmospheric chemistry and physics: from air pollution to climate change*. John Wiley & Sons; 2016 Apr 4.
27. Coccia M. The effects of atmospheric stability with low wind speed and of air pollution on the accelerated transmission dynamics of COVID-19. *International Journal*

of Environmental Studies. 2021 Jan 2;78(1):1-27.