

## An analysis of air pollution trends in Jaipur, UNESCO world heritage city

Mohit Jangir<sup>1</sup>, Parag Jyoti Kashyap<sup>2,\*</sup>, Kheraj<sup>3</sup>, M.P. Punia<sup>4</sup>, Sanchali Das Podder<sup>5</sup>

<sup>1</sup> Government Senior Secondary School Lambi Ahir, Department of School Education Rajasthan, India

<sup>2</sup> Department of Geography, Diphu Campus, Assam University, India

<sup>3</sup> Department of Geography, Central University of Haryana, Haryana, India

<sup>4</sup> Department of Remote Sensing, Birla Institute of Scientific Research, Jaipur, India

<sup>5</sup> Department of Geography, Assam University, Assam, India

### ARTICLE INFORMATION

*Article Chronology:*

Received 07 April 2024

Revised 30 November 2024

Accepted 20 February 2025

Published 29 March 2025

*Keywords:*

Air pollution; Jaipur city; Pollutants;  
Geographic information system (GIS)

### CORRESPONDING AUTHOR:

[jangirmohit97@gmail.com](mailto:jangirmohit97@gmail.com)

Tel : (+91) 8135807474

Fax : (+91) 8135807474

### ABSTRACT

**Introduction:** Introduction: Air pollution is a significant environmental challenge globally, exacerbated by industrialization and increasing vehicular emissions. This study focuses on Jaipur, India, where rapid urbanization and industrial growth have intensified pollution levels, impacting public health and environmental quality.

**Materials and methods:** This study utilized secondary data from the Rajasthan State Pollution Control Board and satellite imagery obtained from the NRSC BHUVAN. Geographic Information System (GIS) tools were employed to analyze pollution data from six sample sites in Jaipur. Interpolation techniques, including Kriging and Inverse Distance Weighting (IDW), were used to map the spatial distribution of pollutants.

**Results:** From 2011 to 2019, Jaipur experienced varying levels of air pollution, with high concentrations of Particulate Matter (PM<sub>10</sub>), Sulfur dioxide (SO<sub>2</sub>), and Nitrogen dioxide (NO<sub>2</sub>) observed in industrial and commercial zones, such as the Vishwakarma Industrial Area and Ajmeri Gate. Areas with natural features, like Jhalana Dungri and the Malaviya Industrial Area, consistently showed lower pollution levels.

**Conclusion:** The study highlights significant spatial and temporal variations in air quality across Jaipur, influenced by industrial activities and vehicular emissions. Effective pollution control measures and urban planning strategies are essential to mitigate the adverse impacts of air pollution on public health and environmental sustainability in Jaipur and similar urban centers.

### Introduction

Environmental degradation has become a major global problem, particularly concerning

air pollution. Since the Industrial Revolution, air pollution has increased rapidly due to large industries emitting substantial amounts of carbon dioxide and suspended particles into the atmosphere as byproducts of burning fossil

Please cite this article as: Jangir M, Kashyap PJ, Kheraj, Punia M.P, Podder SD. An analysis of air pollution trends in Jaipur, UNESCO world heritage city. Journal of Air Pollution and Health. 2025;10(1): 133-154.

fuels [1, 2]. Additionally, industries release many harmful gases and aerosols. Recently, vehicular pollution has escalated with the rising number of vehicles. It is evident that pollution is a significant issue globally [3, 4]. According to a World Health Organization (WHO) report, seven million people die annually due to air pollution, affecting 9 out of 10 people worldwide. All countries are working to control air pollution and improve air quality under the guidance of the World Health Organization. Environmental pollution is a common issue in both developed and developing countries. Each year, substantial quantities of solid waste are released into the atmosphere through the burning of fossil fuels, industrial and domestic waste, and vehicle emissions. In developing nations, urban areas face an environmental emergency primarily due to vehicular emissions, which contribute to 40-80% of total air pollution [5, 6].

In India, air pollution, especially in the northern region, has become a critical issue. The country has 23 significant urban areas with populations exceeding one million, many of which do not meet WHO air quality standards [5, 7]. According to a study, industry contributes 51% to air pollution, vehicle emissions 22%, biomass burning from crops 8%, and other activities-such as household waste, construction, dust storms, and fireworks during festivals-contribute 5%. Air pollution has become a subject of serious discussion due to the rapid increases in human activity in India. Industrialization has led to a daily increase in air pollution, driven by population growth, rising vehicle numbers, biofuel consumption, poor land use patterns, and inadequate infrastructure management [8]. A recent study, "Early Life Exposure to Outdoor Air Pollution: Effects on Child Health in India," found that Particulate Matters ( $PM_{2.5}$ ) affects fetal development, reducing fetal length by 7.9% and weight by 6.7% in the first three months [9].

In India, the main air pollutants are Sulfur

dioxide ( $SO_2$ ), Nitrogen dioxide ( $NO_2$ ), and Suspended Particulate Matter (SPM) [10, 11]. Urban areas are particularly exposed to high levels of air pollution, including metals, due to vehicle emissions, which are the primary source of fine and ultrafine particles [12, 13]. Diesel and petrol fumes from fossil fuels contain harmful substances [14].

Like many Indian cities, Jaipur is facing significant challenges, including urbanization, industrial emissions, traffic congestion, and poor road conditions [15]. These issues, along with its appeal as a tourist destination, have contributed to increased air pollution levels. Similar to other rapidly expanding cities in the nation, Jaipur is experiencing heightened urban development, traffic problems, inadequate infrastructure, and weak regulations on industrial emissions, all of which have adversely affected air quality [16]. While air pollution in the city originates from various sources, a comprehensive analysis of the chemical composition of  $PM_{10}$  and  $PM_{2.5}$  has not yet been performed to identify these sources and their impacts [17]. This emphasizes the critical need to evaluate both gaseous and particulate pollutants over time and across different areas to shape policies and management strategies aimed at reducing air pollution in Jaipur.

### ***Air quality index (AQI)***

The Air Quality Index (AQI) is a daily measure that indicates air quality and its impact on health over a short period. It helps people understand how ambient air quality affects their health [18]. The Environmental Protection Agency (EPA) of the United States calculates the AQI for five common air pollutants, based on national air quality standards designed to protect public health:  $SO_2$ ,  $NO_2$ , CO,  $O_3$ , and PMs (2.5 and 10) [19-21]. Categories of air quality index are shown in Table 1.

Different countries use various methods to calculate air quality; India employs a 500-point scale for its AQI.

Table 1. Categories of air quality index

Value	Type
0-50	Good
51-100	Satisfactory
101-200	Moderately
201-300	Poor
301-400	Very Poor
401-500	Severe

### Objectives

- \* To assess distribution of  $\text{NO}_2$ ,  $\text{PM}_{10}$ ,  $\text{SO}_2$  concentration in Jaipur city
- \* To estimate air quality index of Jaipur city

### Materials and methods

This study relies on secondary data collected from various sources. Satellite imagery of Jaipur city was obtained from NRSC BHUVAN, while pollution data was sourced from the Rajasthan State Pollution Control Board (RSPCB) in Jaipur, Rajasthan. The study encompasses six sample sites designated by the RSPCB to assess pollution levels across different categories, such as industrial and commercial (see Table 2). Data collection followed a standardized procedure, involving the calculation of the annual arithmetic mean based on a minimum of 104 measurements per year at each site, taken twice a week at 24-h intervals. Upon collection, the raw data provided by the pollution control board were classified and

converted into point data. Geographic coordinates were assigned using the ARC Map Tool. The satellite imagery obtained from NRSC BHUVAN was processed into SHP (Shapefile) format and incorporated into the ARC Map Tool, ensuring that all data elements were standardized to a consistent projection. Geographic Information System (GIS) analysis was performed using ARC Map Tool version 10.4. Point data and SHP files were integrated into the tool and aligned with a uniform projection for spatial analysis. The point data were further processed for export, and interpolation techniques such as Kriging and Inverse Distance Weighting (IDW) were applied to spatially distribute air pollution across Jaipur city. Kriging is an interpolation technique that predicts the value of a variable at an unmeasured site based on values from nearby locations [22]. IDW, on the other hand, produces more accurate estimates of PM by weighting measurement data based on distance [23]. After interpolation, the resultant data were formatted into map layouts for visualization and analysis. Methodological flowchart is shown in Fig. 1.

Table 2. Air quality monitoring sample sites of Jaipur city

Sl. No.	Sample sites	Categories
1	Vishwakarma	Industrial area
2	Malaviya	Industrial area
3	Regional Office	Commercial area
4	Jhalana Dungri	Commercial area
5	Ajmeri Gate	Commercial area
6	Chandpole	Commercial area




Fig. 1. Methodological flowchart

### Study area

Geographically located in the northwestern part of India, Rajasthan extends between latitudes  $23^{\circ} 30' N$  and  $30^{\circ} 11' N$ , and longitudes  $69^{\circ} 29' E$  and  $78^{\circ} 17' E$ . Jaipur, the capital and historic city of Rajasthan, is situated in Northern India. According to the 2011 census, Jaipur city has a total population of 3.1 million, making it the 10<sup>th</sup> most populous city in India. Known as the "Pink City" due to the predominant pink color

of its buildings, Jaipur is located approximately 268 km from New Delhi. It is a prominent tourist destination and forms a key part of the Golden Triangle tourist circuit, along with Delhi and Agra (240 km away). The city boasts two United Nations Educational, Scientific and Cultural Organization (UNESCO) world heritage sites: the Amer Fort and the Jantar Mantar Observatory [24, 25]. Location map of the study area is shown in Fig. 2.










Fig. 2. Location map of the study area

**Results and discussion**

Metropolitan air quality is a critical factor influencing public health and economic productivity. Understanding air quality levels, pollutant concentrations, and associated health risks is essential. Urban air pollution has become increasingly complex due to the multitude and diversity of pollution sources, which significantly impact urban air quality. Indian urban areas typically exceed air quality standards and share similarities in development, infrastructure, industrial activities, and commerce with the study area under examination. Various activities, such as cooking and heating with solid fuels, biomass, kerosene, wood, and fossil fuels in open fires or traditional stoves, contribute to high levels of indoor air pollution.

Jaipur, located in northern India, experiences a dry climate and hosts a mix of industrial and commercial activities. The city's ambient air quality is monitored regularly at selected sites across the urban area. This study investigates the spatial and temporal changes in air pollution in Jaipur city from 2011 to 2019. Satellite imagery captured by the LISS 3 satellite was obtained from the NRSC Bhuvan platform for analysis. Interpolation techniques were employed to estimate air pollutant concentrations across the study area. Both Kriging and Inverse Distance Weighted (IDW) methods were utilized for interpolation. IDW differs from Kriging in that it does not involve measurable models or account for spatial autocorrelation. Instead, it uses z-values and distance weights to estimate values in unmeasured areas.











Fig. 3. Air quality index (AQI) in Jaipur city from 2011 to 2019





**Distribution of AQI in Jaipur city**





The maps illustrate the distribution of the Air Quality Index (AQI) in Jaipur city from 2011 to 2019, showing consistently high concentrations in the central part of the city. Specifically, four sample sites—Vishwakarma Industrial Area, the Regional Office, Ajmeri Gate, and Chandpole—are located in this area. Air Quality Index (AQI) in Jaipur city from 2011 to 2019 is shown in Fig. 3.

In 2011, the highest AQI concentrations were recorded in Vishwakarma Industrial Area (197) and the Regional Office (161). By 2012, the highest AQI values shifted to the Regional Office (221) and Ajmeri Gate (192). In subsequent years (2013-2014), AQI peaks returned to Vishwakarma Industrial Area and Ajmeri Gate, reaching 245 and 208, respectively, in 2013, and 205 and 190 in 2014. From 2015 to 2017, the highest AQI concentrations persisted in the Regional Office and Ajmeri Gate, with values ranging from 168

to 231. Additionally, a sudden rise in AQI was observed in the Chandpole area in 2016 (202) and 2017 (175).

In 2018 and 2019, the highest AQI concentrations were again observed in Vishwakarma Industrial Area, the Regional Office, the Regional Office, and Ajmeri Gate, with values varying from 140 to 234. These high concentrations are attributed to industrial activities, commercial operations, and heavy traffic in these areas. Throughout the period from 2011 to 2019, the lowest AQI concentrations were consistently found in Jhalana Doongri and Malaviya Industrial Area. PM<sub>10</sub> values in Jhalana Doongri ranged from 90 to 120, while in Malaviya Industrial Area; they ranged from 100 to 119. These areas benefit from natural features such as forests (Jhalana Doongri and Nahargarh Forest Reserve) and the hilly terrain of the Aravalli Mountains, resulting in relatively lower AQI concentrations.













Fig. 4. PM<sub>10</sub> levels in Jaipur city from 2011 to 2019





These maps depict the distribution of  $PM_{10}$  pollution in Jaipur city from 2011 to 2019, showing significantly high concentrations in the northern and central parts, where four sample sites, Vishwakarma industrial area (VKI), the Regional Office (RO), Ajmeri Gate, and Chandpole are located. The highest  $PM_{10}$  concentrations were consistently found in Vishwakarma Industrial Area, Ajmeri Gate, and the Regional Office. Fig. 4 shows the  $PM_{10}$  levels in Jaipur city from 2011 to 2019.

In 2011,  $PM_{10}$  levels were  $216 \mu\text{g}/\text{m}^3$  at Ajmeri Gate and  $246 \mu\text{g}/\text{m}^3$  at Vishwakarma Industrial Area. By 2012, Ajmeri Gate recorded  $238 \mu\text{g}/\text{m}^3$ , Vishwakarma Industrial Area had  $163 \mu\text{g}/\text{m}^3$ , and the Regional Office reached  $271 \mu\text{g}/\text{m}^3$ . In subsequent years,  $PM_{10}$  concentrations fluctuated: in 2013, Ajmeri Gate had  $258 \mu\text{g}/\text{m}^3$ , Vishwakarma industrial area  $295 \mu\text{g}/\text{m}^3$ , and the Regional Office  $197 \mu\text{g}/\text{m}^3$ ; in 2014, Ajmeri Gate recorded  $236 \mu\text{g}/\text{m}^3$ , Vishwakarma industrial area  $255 \mu\text{g}/\text{m}^3$ , and the Regional Office  $225 \mu\text{g}/\text{m}^3$ ; in 2015, Ajmeri Gate had  $249 \mu\text{g}/\text{m}^3$ , Vishwakarma Industrial Area  $201 \mu\text{g}/\text{m}^3$ , Chandpole  $256 \mu\text{g}/\text{m}^3$ , and the Regional Office  $230 \mu\text{g}/\text{m}^3$ ; in 2016,

Ajmeri Gate reached  $281 \mu\text{g}/\text{m}^3$ , Vishwakarma Industrial Area  $159 \mu\text{g}/\text{m}^3$ , Chandpole  $213 \mu\text{g}/\text{m}^3$ , and the Regional Office  $252 \mu\text{g}/\text{m}^3$ . From 2017 to 2019,  $PM_{10}$  concentrations remained high, with Ajmeri Gate recording  $272 \mu\text{g}/\text{m}^3$ ,  $284 \mu\text{g}/\text{m}^3$ , and  $193 \mu\text{g}/\text{m}^3$ ; Vishwakarma Industrial Area with  $159 \mu\text{g}/\text{m}^3$ ,  $207 \mu\text{g}/\text{m}^3$ , and  $207 \mu\text{g}/\text{m}^3$ ; and the Regional Office with  $251 \mu\text{g}/\text{m}^3$ ,  $151 \mu\text{g}/\text{m}^3$ , and  $202 \mu\text{g}/\text{m}^3$ . These areas are characterized by heavy commercial and residential activities, contributing to significant traffic congestion and heightened  $PM_{10}$  levels, which exceed the permissible limit of  $60 \mu\text{g}/\text{m}^3$ .

In contrast, the southern parts of Jaipur city showed moderate  $PM_{10}$  concentrations, while consistently lower levels were observed in Jhalana Dungri and Malaviya Industrial Area throughout 2011 to 2019. Jhalana Dungri recorded concentrations ranging from  $90 \mu\text{g}/\text{m}^3$  to  $130 \mu\text{g}/\text{m}^3$ , and Malaviya Industrial Area from  $103 \mu\text{g}/\text{m}^3$  to  $129 \mu\text{g}/\text{m}^3$ . These areas benefit from natural features such as forests (Jhalana Dungri and Nahargarh Forest Reserve) and the hills of the Aravalli Mountains, resulting in comparatively lower  $PM_{10}$  levels.














Fig. 5. SO<sub>2</sub> levels in Jaipur city from 2011 to 2019

SO<sub>2</sub> pollution is notably high in the northern and central parts of Jaipur city, where four sample sites—Vishwakarma Industrial Area (VKI), the Regional Office (RO), Ajmeri Gate, and Chandpole—are located. Throughout the years, the highest concentrations of SO<sub>2</sub> were consistently observed in Vishwakarma Industrial Area, Ajmeri Gate, Chandpole, and the Regional Office. From 2011 to 2019, SO<sub>2</sub> concentrations in Vishwakarma Industrial Area ranged from 12 µg/m<sup>3</sup> to 13 µg/m<sup>3</sup>, in Chandpole from 6 µg/m<sup>3</sup> to 9 µg/m<sup>3</sup>, in Ajmeri Gate from 6 µg/m<sup>3</sup> to 10 µg/m<sup>3</sup>, and in the Regional Office from 7 µg/m<sup>3</sup> to 10 µg/m<sup>3</sup>. These areas are characterized by heavy industrial and commercial activities,

contributing to elevated levels of SO<sub>2</sub> pollution. Fig. 5 shows SO<sub>2</sub> levels in Jaipur city from 2011 to 2019.

Conversely, the lowest SO<sub>2</sub> concentrations were consistently found in Jhalana Doongri and Malaviya Industrial Area throughout the same period. In Jhalana Doongri, concentrations ranged from 6 µg/m<sup>3</sup> to 8 µg/m<sup>3</sup>, and in Malaviya Industrial Area from 6 µg/m<sup>3</sup> to 9 µg/m<sup>3</sup>. These areas benefit from natural features such as forests (Jhalana Doongri and Nahargarh Forest Reserve) and the hilly terrain of the Aravalli Mountains, resulting in comparatively lower SO<sub>2</sub> levels. The permissible limit for SO<sub>2</sub> is 50 µg/m<sup>3</sup>, which was not exceeded at any of the sample sites in 2011.










Fig. 6. NO<sub>2</sub> levels in Jaipur city from 2011 to 2019

NO<sub>2</sub> concentrations varied across different areas of Jaipur city from 2011 to 2019. The highest concentrations were observed in Chandpole in 2011 and during 2014-2015, in the Regional Office in 2012-2013 and 2016, and in Ajmeri Gate in 2016-2018. By 2019, the highest NO<sub>2</sub> levels were found in Vishwakarma Industrial Area. Specifically, in Ajmeri Gate, NO<sub>2</sub> concentrations were 38 µg/m<sup>3</sup> in 2011, increasing to 53 µg/m<sup>3</sup> in 2012, and varying between 31 µg/m<sup>3</sup> and 37 µg/m<sup>3</sup> from 2013 to 2019. Vishwakarma Industrial Area recorded levels ranging from 21 µg/m<sup>3</sup> to 37 µg/m<sup>3</sup> over the same period. Chandpole saw concentrations from 39 µg/m<sup>3</sup> to 52 µg/m<sup>3</sup>, while the Regional Office ranged from 27 µg/m<sup>3</sup> to 54 µg/m<sup>3</sup>. Conversely, the lowest NO<sub>2</sub> concentrations were consistently found in Jhalana Doongri and Malaviya Industrial Area throughout these years. In Jhalana Doongri, concentrations ranged from 21 µg/m<sup>3</sup> to 46 µg/m<sup>3</sup>, and in Malaviya Industrial Area from 21 µg/m<sup>3</sup> to 47 µg/m<sup>3</sup>. These areas benefit from natural features and have comparatively lower NO<sub>2</sub> levels due to reduced industrial and traffic activities. Fig. 6 shows NO<sub>2</sub> levels in Jaipur city from 2011 to 2019.

The analysis of air quality in Jaipur from 2011 to 2019 highlights a complex interplay between urban expansion, industrial activities, and environmental factors that significantly impact public health and economic outcomes. The results reveal considerable challenges with air pollution, especially in urban areas where pollution levels frequently surpass national standards. Understanding these dynamics is crucial for developing effective public policies and urban planning strategies to mitigate the health impacts of pollution. The data indicates a distinct contrast in air quality between industrial and natural areas in Jaipur. Regions such as the Vishwakarma Industrial Area and Ajmeri Gate consistently report high Air Quality Index (AQI) values due to emissions from factories, vehicles, and construction. In contrast, residential or commercial areas like

Jhalana Doongri area show significantly lower AQI and particulate matter (PM<sub>10</sub>) levels, benefiting from their natural surroundings. This disparity highlights the need for targeted interventions in the most polluted regions. Throughout the study period, pollutants including PM<sub>10</sub>, sulfur dioxide (SO<sub>2</sub>), and nitrogen dioxide (NO<sub>2</sub>) displayed troubling trends, with the highest concentrations located in the central and northern parts of the city. PM<sub>10</sub> levels often exceeded the permissible limit of 60 µg/m<sup>3</sup>, particularly in the Vishwakarma Industrial Area, where they peaked at 295 µg/m<sup>3</sup> in 2013. This ongoing pollution reflects the high level of industrial and commercial activities in these areas, posing serious health risks to local residents. Conversely, the lower pollution levels in Jhalana Doongri area is likely due to natural barriers and vegetation that help filter pollutants. These findings suggest that integrating green spaces into urban planning could significantly improve air quality. Several interconnected factors contribute to the observed air quality trends in Jaipur. Rapid urbanization and population growth have driven increased industrial output and emissions. Traditional practices, such as using solid fuels for cooking and heating in densely populated areas, worsen indoor air pollution. The dependence on fossil fuels for transportation exacerbates urban pollution, with vehicle emissions being a major contributor. Furthermore, inadequate waste management and ongoing construction activities elevate pollution levels, as dust from construction sites worsens air quality. Addressing these challenges requires a comprehensive strategy that incorporates both environmental and socio-economic considerations. The health implications of persistently high pollution levels in Jaipur are severe. Exposure to elevated PM<sub>10</sub> and other pollutants is linked to various health issues, including respiratory and cardiovascular diseases. Vulnerable populations, particularly children and the elderly, face the greatest risks, highlighting the urgent need for

effective public health measures. Continuous monitoring of health outcomes related to air quality is essential for understanding the full impact of pollution on community health. Public awareness campaigns that inform residents about the risks of air pollution and promote healthier behaviors can significantly help reduce health threats. Encouraging the use of cleaner cooking fuels and improving waste disposal methods can greatly decrease both indoor and outdoor pollution levels.

Similar to Jaipur, other World Heritage cities in India, including Chandigarh and Ahmedabad, are facing serious air pollution challenges. In Chandigarh, the pollution statistics present a mixed picture: PM<sub>10</sub> levels have decreased at industrial sites, but NO<sub>x</sub> levels are increasing throughout the city. Seasonal changes significantly impact pollutant levels, often resulting in higher concentrations during winter and after the monsoon, primarily due to construction and dust. Although there have been some positive developments, air quality remains a major concern, especially for vulnerable populations experiencing heightened respiratory issues [26, 27]. In Ahmedabad, some areas have shown improvements, with SO<sub>2</sub> and NO<sub>2</sub> levels consistently below national standards. However, there is a concerning rise in Respirable Suspended Particulate Matter (RSPM) and Suspended Particulate Matter (SPM) at specific monitoring stations, largely due to rapid urbanization, population growth, and an increase in vehicles. While efforts to manage certain pollutants have achieved some success, overall air quality still poses health risks, emphasizing the need for ongoing monitoring and intervention [28, 29].

## **Conclusion**

This study provides a comprehensive assessment of pollution levels in Jaipur city, revealing higher concentrations in the northern

parts compared to lower levels in the southern areas. These findings were determined using interpolation techniques in ArcGIS 10.4, based on pollution data from various locations. The study found that PM<sub>10</sub> pollutants formed a predominant cluster with higher concentrations than vapor pollutants. Pollution concentrations varied significantly across different areas due to local activities. The northern part of Jaipur, characterized by industrial establishments, exhibited the highest pollution levels. In the central areas, where commercial and residential activities overlap, pollutants from non-renewable energy sources and heavy traffic contributed to similarly high concentrations. Conversely, pollution levels were notably lower in areas like Jhalana Doongri and the Malaviya Industrial Area (MIA), which can be attributed to ample forest cover and the nature of industries operating there. Historically, the Vishwakarma Industrial Area (VKI) has been one of the most polluted zones in Jaipur due to its concentration of major industries, with pollution levels showing an upward trajectory over the years. The primary air quality concerns in Jaipur revolve around the Air Quality Index (AQI) and PM<sub>10</sub> levels. SO<sub>2</sub> and NO<sub>2</sub> pollutants were found to be within permissible limits, with NO<sub>2</sub> emissions sourced from activities such as burning fossil fuels, coal, and industrial processes like manufacturing and food processing. SO<sub>2</sub> primarily originates from industrial emissions. Sources of PM<sub>10</sub> include brickworks, refineries, cement and iron industries, fossil fuel combustion, sandstorms, fires, and certain consumer products.

The study emphasizes the importance of a proactive approach to urban planning and public policy in Jaipur. Policymakers should prioritize environmental sustainability alongside economic growth by implementing stricter regulations on industrial emissions, promoting cleaner technologies, and enhancing public transportation to reduce reliance on private vehicles. Additionally, incorporating

green spaces into urban design can enhance air quality and improve residents' overall quality of life. Collaboration among government agencies, industries, and the community is crucial for developing effective policies to address air quality challenges. Ongoing research and data collection will be vital for informing these policies and ensuring that interventions adapt to the changing landscape of urban air pollution.

### Financial supports

No financial support was received for this study.

### Competing interests

No conflicts of interest exist. The study aimed to assess air pollution levels in Jaipur city of India from 2011 to 2019.

### Authors' contributions

Mohit Jangir conducted the study under the supervision of Dr. Kheraj and Dr. M.P. Punia.

Mohit Jangir: Contributed to data compilation, interpretation of results, and writing of the research paper.

Parag Jyoti Kashyap: Contributed to data compilation and manuscript preparation.

Sanchali Das Podder: Edited and arranged the writing.

### Acknowledgements

Data were acquired from the Rajasthan State Pollution Control Board. The authors gratefully acknowledge the Rajasthan State Pollution Control Board for providing the data. We would also like to extend our sincere thanks to Dr. Neha Pandey and Dr. Kutub Beg from the Department of Remote Sensing at BISR Jaipur for their invaluable support throughout the study.

### Ethical considerations

“Ethical issues (Including plagiarism, Informed Consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc) have been completely observed by the authors.”

### References

1. Ghose MK, Paul R, Banerjee SK. Assessment of the impacts of vehicular emissions on urban air quality and its management in Indian context: The case of Kolkata (Calcutta). *Environ Sci Policy*. 2004 Jul; 345-351.
2. Kumar K, Pande BP. Air pollution prediction with machine learning: a case study of Indian cities. *Int J Environ Sci Technol*. 2023;20(5):5333-5348.
3. Gopu P, Panda RR, Nagwani NK. Time series analysis using ARIMA model for air pollution prediction in Hyderabad city of India. In: *Soft Computing and Signal Processing: Proceedings of 3rd ICSCSP 2020, Volume 1*. Singapore: Springer; 2021; p. 47-56.
4. Tiwari N, Kashyap PJ. Urban growth dynamics of National Capital Region of India using geospatial technology. *Asian J Geogr Res*. 2024;7(2):35-52.
5. Gupta HK, Gupta VB, Rao CVC, Gajghate DG, Hasan MZ. Urban ambient air quality and its management strategy for a metropolitan city in India. *Bull Environ Contam Toxicol*. 2002;68(3).
6. Agrawal G, Mohan D, Rahman H. Ambient air pollution in selected small cities in India: Observed trends and future challenges. *IATSS Res*. 2021;45(1):19-30.
7. Malekafzali S, Jozi SA, Kashefiolasl M, Zarimdar M, Shati M. Scientometric analysis of health impact assessment of outdoor air pollution by WHO-AirQ tool (2005-2019). *J Air Pollut Health*. 2021;6(2):101-116.
8. Sicard P, Agathokleous E, Anenberg SC, De

- Marco A, Paoletti E, Calatayud V. Trends in urban air pollution over the last two decades: A global perspective. *Sci Total Environ.* 2023; 858:160064.
9. Singh P, Dey S, Chowdhury S, Bali K. Early life exposure to outdoor air pollution: effect on child health in India. 2019.
10. Agrawal M, Singh J. Impact of coal power plant emission on the foliar elemental concentrations in plants in a low rainfall tropical region. *Environ Monit Assess.* 2000; 60:261-282.
11. Sharma D, Mauzerall D. Analysis of air pollution data in India between 2015 and 2019. *Aerosol Air Qual Res.* 2022; 22(2):210204.
12. Sharma K, Singh R, Barman SC, Mishra D, Kumar R, Negi MPS, et al. Comparison of trace metals concentration in PM<sub>10</sub> of different locations of Lucknow City, India. *Bull Environ Contam Toxicol.* 2006; 77(3):419-426.
13. Gulia S, Shukla N, Padhi L, Bosu P, Goyal SK, Kumar R. Evolution of air pollution management policies and related research in India. *Environ Challenges.* 2022;6:100431.
14. Gurjar BR. Air pollution in India: Major issues and challenges. *Energy Future.* 2021.
15. Ruhela M, Maheshwari V, Ahamad F, Kamboj V. Air quality assessment of Jaipur city Rajasthan after the COVID-19 lockdown. *Spatial Information Research.* 2022 Oct;30(5):597-605.
16. Dadhich AP, Goyal R, Dadhich PN. Assessment of spatio-temporal variations in air quality of Jaipur city, Rajasthan, India. *The Egyptian journal of remote sensing and space science.* 2018 Sep 1;21(2):173-81.
17. Jain S, Mandowara VL. Study on particulate matter pollution in jaipur city. *International Journal of Applied Engineering Research.* 2019;14(3):637-45.
18. Manisalidis I, Stavropoulou E, Stavropoulos A, Bezirtzoglou E. Environmental and health impacts of air pollution: a review. *Front Public Health.* 2020;8:14.
19. Sicard P, Agathokleous E, Anenberg SC, De Marco A, Paoletti E, Calatayud V. Trends in urban air pollution over the last two decades: A global perspective. *Sci Total Environ.* 2023;858:160064.
20. Pathak M, Kuttippurath J. Air quality trends in rural India: analysis of NO<sub>2</sub> pollution using satellite measurements. *Environ Sci Process Impacts.* 2022;24(12):2437-2449.
21. Kooranifar S, Attar GA, Talebi A, Pourashraf M, Rostami R. Short-term effects of exposure to PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, CO, SO<sub>2</sub>, O<sub>3</sub> on lung function test indices among students of Tehran city, Iran. *J Air Pollut Health.* 2021.
22. Liu KF, Hung MJ, Kuo JY, Liang HH. Using GIS and kriging to analyze the spatial distributions of the health risk of indoor air pollution. *Journal of Geoscience and Environment Protection.* 2015 Aug 25;3(6):20-5.
23. Choi K, Chong K. Modified inverse distance weighting interpolation for particulate matter estimation and mapping. *Atmosphere.* 2022 May 22;13(5):846.
24. Jawaid MF, Sharma M, Pipralia S, Kumar A. City profile: Jaipur. *Cities.* 2017;68:63-81.
25. Doshi BV. City of Jaipur. *Architecture Plus Design.* 1989;5(2):96.
26. Saharan S, Deswal S, Pal M. Air Quality Mapping and Urban Planning for Sustainable Urban Ecology: A Case Study of Chandigarh, India. *Ecological Questions.* 2024;35(2):1-5.
27. Ravindra K, Singh T, Pandey V, Mor S. Air pollution trend in Chandigarh city situated in Indo-Gangetic Plains: Understanding seasonality and impact of mitigation strategies. *Science of the Total Environment.* 2020 Aug 10;729:138717.
28. Pathak CY, Mandalia HC, Roy D, Jadeja RB. Comparative study of ambient air quality

status of Ahmedabad and Gandhinagar City in Gujarat, India. *Chemical Science Transactions*. 2015;4(1):89-94.

29. Maltare NN, Vahora S. Air Quality Index prediction using machine learning for Ahmedabad city. *Digital Chemical Engineering*. 2023 Jun 1;7:100093.